

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 March 2006 (09.03.2006)

PCT

(10) International Publication Number
WO 2006/025783 A1

(51) International Patent Classification⁷: **C07D 241/26**,
241/14, 241/12, 213/24, 213/04, 403/04, 401/04, 333/32,
327/04, 231/14, C07C 233/65, A61K 31/192, 31/166,
31/4965, 31/497, 31/435, A61P 9/10, 11/00, 19/02, 37/00

WILLIS, Paul [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough Leicestershire LE11 5RH (GB).

(21) International Application Number:
PCT/SE2005/001251

(74) Agent: **ASTRAZENECA**; Global Intellectual Property, S-151 85 Södertälje (SE).

(22) International Filing Date: 29 August 2005 (29.08.2005)

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0402103-6 30 August 2004 (30.08.2004) SE
0403054-0 15 December 2004 (15.12.2004) SE
0500766-1 6 April 2005 (06.04.2005) SE

(71) Applicant (*for all designated States except US*): **ASTRAZENECA AB** [SE/SE]; S-151 85 Södertälje (SE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FORD, Rhonan** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough Leicestershire LE11 5RH (GB). **MARTIN, Barrie** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough Leicestershire LE11 5RH (GB). **THOMPSON, Toby** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough Leicestershire LE11 5RH (GB). **TOMKINSON, Nicholas** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough Leicestershire LE11 5RH (GB).


(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ADAMANTYL DERIVATES AS P2X7 RECEPTOR ANTAGONISTS



(I)

(57) Abstract: The invention provides compounds of formula (I) pharmaceutically acceptable salt or solvate thereof, in which R¹, A¹, m and A are as defined in the specification; a process for their preparation; pharmaceutical compositions containing them; and their use in therapy.

WO 2006/025783 A1

Adamantyl derivatives as P2X₇ receptor antagonists

The present invention relates to adamantyl derivatives, a process for their preparation,
5 pharmaceutical compositions containing them, a process for preparing pharmaceutical compositions and their use in therapy.


The P2X₇ receptor (previously known as P2Z receptor), which is a ligand-gated ion
channel, is present on a variety of cell types, largely those known to be involved in the
10 inflammatory/immune process, specifically, macrophages, mast cells and lymphocytes
(T and B). Activation of the P2X₇ receptor by extracellular nucleotides, in particular
adenosine triphosphate, leads to the release of interleukin-1 β (IL-1 β) and giant cell
formation (macrophages/microglial cells), degranulation (mast cells) and proliferation
(T cells), apoptosis and L-selectin shedding (lymphocytes). P2X₇ receptors are also
15 located on antigen-presenting cells (APC), keratinocytes, salivary acinar cells (parotid
cells), hepatocytes and mesangial cells.

It would be desirable to make compounds effective as P2X₇ receptor antagonists for use in
the treatment of inflammatory, immune or cardiovascular diseases, in the aetiologies of
20 which the P2X₇ receptor may play a role.

The present invention provides a new class of adamantyl-containing P2X₇ antagonist that
comprise a substituted biaromatic group. These novel compounds display excellent
properties for use as P2X₇ receptor antagonists in the treatment of inflammatory, immune
25 or cardiovascular diseases. Whilst adamantyl-containing P2X₇ antagonists have been
described previously, for example in WO 00/61569, WO 03/080579 and WO 03/042190,
prior to the present invention there had been no suggestion that compounds comprising the
substituted biaromatic group of the present invention would make good P2X₇ antagonists.
US patent application 2003/0134885 A1 concerns substituted biphenyl ligand activators of

PPARgamma receptors. It does not mention the P2X₇ receptor or describe any adamantyl derivatives.

In accordance with the present invention, there is therefore provided a compound of
 5 general formula (I), or a pharmaceutically acceptable salt or solvate thereof,




wherein m represents 1, 2 or 3;


each R¹ independently represents a hydrogen atom or a halogen;

A represents C(O)NH or NHC(O);


10 Ar¹ represents a group



(II)




(III)



(IV)

or



(V)

15 one of R² and R³ represents halogen, nitro, NR⁴R⁵, hydroxyl, or a group selected from (i) C₁-C₆ alkyl optionally substituted by at least one halogen and (ii) C₁-C₆ alkoxy optionally substituted by at least one halogen, and the other of R² and R³ represents a hydrogen atom, halogen or a C₁-C₆ alkyl group optionally substituted by at least one halogen;

20 R⁴ and R⁵ each independently represent a hydrogen atom or a group selected from C₁-C₆ alkyl and C₁-C₆ alkoxy, which C₁-C₆ alkyl or C₁-C₆ alkoxy group can be optionally substituted with at least one substituent selected from halogen and hydroxyl;

Ar² represents phenyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, which phenyl or heteroaromatic ring is substituted by at least one substituent selected from CO₂R⁶, MC₁₋₆ alkylCO₂R⁷, C₁₋₆ alkylsulphonylaminocarbonyl, NHR⁸, R⁹, XR¹⁰, C(O)NHOH and
 5 NR²⁸R²⁹;

and which phenyl or heteroaromatic ring can further be optionally substituted by at least one substituent selected from halogen, nitro, NR¹¹R¹², hydroxyl, S(O)_pR¹³, a C₁-C₆ alkoxy group which C₁-C₆ alkoxy group can be optionally substituted by at least one halogen, and a C₁-C₆ alkyl group which C₁-C₆ alkyl group can be optionally substituted by at least one
 10 substituent selected from halogen, hydroxyl, NR¹⁴R¹⁵, SO₂NR¹⁶R¹⁷, NR¹⁸SO₂R¹⁹, NHCOR²⁰ and CONR²¹R²²;

R⁶ and R⁷ each independently represent a hydrogen atom or a C₁-C₆ alkyl group;

R⁸ represents CN, C₁-C₆ alkylsulphonyl, C₁-C₆ alkylcarbonyl, C₁-C₆ alkoxycarbonyl, C₁-C₆ alkylaminosulphonyl, or (di)-C₁-C₆ alkylaminosulphonyl;

15 R⁹ and R¹⁰ each independently represent tetrazolyl or a 5- to 6-membered heterocyclic ring comprising from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulphur, which heterocyclic ring is substituted by at least one substituent selected from hydroxyl, =O and =S;

M represents a bond, oxygen, S(O)_q or NR²³;

20 X represents oxygen, S(O)_s, NR²⁴, C₁-C₆ alkylene, O(CH₂)₁₋₆, NR²⁵(CH₂)₁₋₆, or S(O)_t(CH₂)₁₋₆;

p, q, s and t each independently represent 0, 1 or 2;

R²⁸ and R²⁹ together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring, which heterocyclic ring is substituted with at least
 25 one substituent independently selected from CO₂R⁶, MC₁₋₆ alkylCO₂R⁷, C₁₋₆ alkylsulphonylaminocarbonyl, C(O)NHOH, NHR⁸, R⁹ and XR¹⁰, and which 3- to 8-membered saturated heterocyclic ring can further be optionally substituted by at least one substituent independently selected from hydroxyl, halogen, C₁-C₆ alkoxy optionally substituted by at least one halogen, and a C₁-C₆ alkyl group which C₁-C₆ alkyl group can
 30 be optionally substituted by at least one substituent independently selected from halogen and hydroxyl; and

R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} and R^{25} each independently represent a hydrogen atom or a group selected from C_1 - C_6 alkyl and C_1 - C_6 alkoxy, which C_1 - C_6 alkyl or C_1 - C_6 alkoxy group can be optionally substituted with at least one substituent selected from halogen and hydroxyl;

provided that:

- when m is 1 and Ar^1 is a group (II) and Ar^2 is phenyl substituted by XR^{10} in a position para to Ar^1 and X is CH_2 , then R^{10} is not a 2,4-dioxothiazolyl group, and
- when m is 1 and Ar^1 is a group (II) and Ar^2 is phenyl substituted by $MC_{1-6}alkylCO_2R^7$ in a position para to Ar^1 , then M does not represent a bond.

Certain compounds of formula (I) are capable of existing in stereoisomeric forms. It will be understood that the invention encompasses all geometric and optical isomers of the compounds of formula (I) and mixtures thereof including racemates. Tautomers and mixtures thereof also form an aspect of the present invention.

In the context of the present specification, unless otherwise indicated, a "Heterocyclic" ring is an unsaturated, saturated or partially saturated ring, at least one atom of which is a heteroatom selected from oxygen, sulphur or nitrogen, and may have aliphatic or aromatic properties. "Heteroaromatic" denotes aromatic rings, at least one atom of which is a heteroatom selected from oxygen, sulphur or nitrogen. A "Carbocyclic" ring is an unsaturated, saturated or partially saturated ring, containing only carbon ring atoms, and may have aliphatic or aromatic properties. The term "Cycloalkyl" denotes saturated alkyl rings. Unless otherwise indicated an alkyl group may be linear or branched. Where a ring or group is described as being optionally substituted with at least one substituent the ring or group may be unsubstituted, or alternatively the ring or group may be substituted with, for example, one, two or three substituents.

In an embodiment of the invention, m represents 1. In another embodiment of the invention, m represents 2.

In an embodiment of the invention, each R^1 independently represents a hydrogen atom.


In an embodiment of the invention, A represents $NHC(O)$.

In an embodiment of the invention, Ar^1 represents a group (II) or (III).

One of R^2 and R^3 represents halogen (e.g. fluorine, chlorine, bromine or iodine), nitro, NR^4R^5 , hydroxyl, or a group selected from (i) $\text{C}_1\text{-C}_6$, preferably $\text{C}_1\text{-C}_4$, alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) optionally substituted by at least one (e.g. zero, one, two or three) halogen (e.g. fluorine, chlorine, bromine or iodine), and (ii) $\text{C}_1\text{-C}_6$, preferably $\text{C}_1\text{-C}_4$, alkoxy (e.g. methoxy, ethoxy, n-propoxy, n-butoxy, n-pentoxy or n-hexoxy) optionally substituted by at least one (e.g. zero, one, two or three) halogen (e.g. fluorine, chlorine, bromine or iodine), and the other of R^2 and R^3 represents a hydrogen atom, halogen (e.g. fluorine, chlorine, bromine or iodine) or a $\text{C}_1\text{-C}_6$, preferably $\text{C}_1\text{-C}_4$, alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) group optionally substituted by at least one (e.g. zero, one, two or three) halogen (e.g. fluorine, chlorine, bromine or iodine).

In an embodiment of the invention, R^2 represents halogen, nitro, NH_2 , hydroxyl, or a $\text{C}_1\text{-C}_6$ alkyl optionally substituted by one to three halogen substituents; and R^3 represents a hydrogen atom.

In an embodiment of the invention, Ar^1 represents a group



wherein R^2 represents halogen (e.g. fluorine, chlorine, bromine or iodine), nitro, NH_2 , hydroxyl, or a C_1 - C_6 , preferably C_1 - C_4 , alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) optionally substituted by at least one (e.g. zero, one, two or three) halogen (e.g. fluorine, chlorine, bromine or iodine).

5

R^4 and R^5 each independently represent a hydrogen atom or a group selected from C_1 - C_6 , preferably C_1 - C_4 , alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) and C_1 - C_6 , preferably C_1 - C_4 , alkoxy (e.g. methoxy, ethoxy, n-propoxy, n-butoxy, n-pentoxy or n-hexoxy), which C_1 - C_6 alkyl or C_1 - C_6 alkoxy group can be optionally substituted with at least one (e.g. zero, one, two or three) substituent selected from halogen (e.g. fluorine, chlorine, bromine or iodine), and hydroxyl.

10

Ar^2 represents phenyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, which phenyl or heteroaromatic ring is substituted by at least one (e.g. one or two) substituent selected from CO_2R^6 , MC_1 - C_6 alkyl CO_2R^7 , C_1 - C_6 alkylsulphonylaminocarbonyl (e.g. $MeSO_2NHCO$ -, or $EtSO_2NHCO$ -), NHR^8 , R^9 , XR^{10} , $C(O)NHOH$ and $NR^{28}R^{29}$; and which phenyl or heteroaromatic ring can further be optionally substituted by at least one (e.g. zero, one or two) substituent selected from halogen (e.g. fluorine, chlorine, bromine or iodine), nitro, $NR^{11}R^{12}$, hydroxyl, $S(O)_pR^{13}$, a C_1 - C_6 , preferably C_1 - C_4 , alkoxy (e.g. methoxy, ethoxy, n-propoxy, n-butoxy, n-pentoxy or n-hexoxy) group which C_1 - C_6 alkoxy group can be optionally substituted by at least one halogen (e.g. fluorine, chlorine, bromine or iodine), and a C_1 - C_6 , preferably C_1 - C_4 , alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) group which C_1 - C_6 alkyl group can be optionally substituted by at least one (e.g. zero, one or two) substituent selected from halogen (e.g. fluorine, chlorine, bromine or iodine), hydroxyl, $NR^{14}R^{15}$, $SO_2NR^{16}R^{17}$, $NR^{18}SO_2R^{19}$, $NHCOR^{20}$ and $CONR^{21}R^{22}$.

25

When Ar^2 represents phenyl or a 6-membered heteroaromatic ring, the at least one substituent selected from CO_2R^6 , MC_1 - C_6 alkyl CO_2R^7 , C_1 - C_6 alkylsulphonylaminocarbonyl,

30

NHR⁸, R⁹, XR¹⁰, C(O)NHOH and NR²⁸R²⁹ may be positioned in an ortho, meta or para position relative to the bond between Ar¹ and Ar². In an embodiment of the invention, when Ar² represents phenyl or a 6-membered heteroaromatic ring the at least one substituent is in an ortho position relative to the bond between Ar¹ and Ar². In another embodiment of the invention, the at least one substituent is in a meta position relative to the bond between Ar¹ and Ar².

Examples of 5- or 6-membered heteroaromatic rings that Ar² may represent include pyrrolyl, thienyl, furanyl, imidazolyl, thiazolyl, oxazolyl, pyrazolyl, isothiazolyl, isoxazolyl, pyridyl, pyridazinyl, pyrimidinyl and pyrazinyl.

In an embodiment of the invention, Ar² represents phenyl, thienyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 nitrogen atoms. In a further embodiment of the invention, Ar² represents phenyl, thienyl or a 6-membered heteroaromatic ring comprising from 1 to 2 nitrogen atoms, e.g. pyridyl, pyridazinyl, pyrimidinyl or pyrazinyl. In another embodiment of the invention, Ar² represents phenyl, thienyl, pyridyl, pyrazolyl or pyrazinyl. In another embodiment of the invention Ar² represents phenyl or pyridyl.

In an embodiment of the invention, Ar² is substituted by at least one (e.g. one or two) substituent selected from carboxyl, -C₁-C₆alkylCO₂H, -OC₁-C₆alkylCO₂H, -NHC₁-C₆alkylCO₂H, -N(C₁-C₄alkyl)C₁-C₆alkylCO₂H, -NHCN, -NHCOC₁-C₆alkyl, -NHSO₂C₁-C₆alkyl, -CONHSO₂C₁-C₆alkyl, tetrazolyl and -OC₁-C₆alkyltetrazolyl. In a further aspect of this embodiment Ar² can further be optionally substituted by at least one substituent (e.g. zero, one or two) selected from halogen, trifluoromethyl, C₁-C₆alkoxy and a C₁-C₆alkyl group.

In another embodiment of the invention, Ar² is substituted by a group NR²⁸R²⁹, wherein R²⁸ and R²⁹ together with the nitrogen atom to which they are attached form a saturated heterocyclic group selected from azetidiny, pyrrolidinyl or piperidinyl, which heterocyclic group is substituted by carboxyl and can further be optionally substituted by hydroxyl.

In a further embodiment of the invention Ar^2 is substituted by a substituent selected from carboxyl, MC_1-C_6 alkyl CO_2R^7 and C_1-C_6 alkylsulphonylaminocarbonyl, and can further be optionally substituted by at least one substituent selected from halogen and a C_1-C_6 alkyl group.

5

In another embodiment of the invention Ar^2 is substituted by carboxyl, and optionally at least one (e.g. zero, one or two) further substituent selected from halogen and a C_1-C_6 alkyl group.

10

R^6 and R^7 each independently represent a hydrogen atom or a C_1-C_6 , preferably C_1-C_4 , alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) group. In an embodiment of the invention, R^6 and R^7 each independently represent a hydrogen atom.

15

R^8 represents CN, C_1-C_6 , preferably C_1-C_4 , alkylsulphonyl (e.g. $MeSO_2-$ or $EtSO_2-$), C_1-C_6 , preferably C_1-C_4 , alkylcarbonyl (e.g. methyl-, ethyl-, n-propyl-, n-butyl-, n-pentyl- or n-hexylcarbonyl), C_1-C_6 , preferably C_1-C_4 , alkoxycarbonyl (e.g. methoxy-, ethoxy-, n-propoxy-, n-butoxy-, n-pentoxo- or n-hexoxycarbonyl), C_1-C_6 , preferably C_1-C_4 , alkylaminosulphonyl (e.g. $MeNHSO_2$ or $EtNHSO_2-$), or (di)- C_1-C_6 , preferably C_1-C_4 ,
20 alkylaminosulphonyl (e.g. Me_2NSO_2 or Et_2NSO_2- or $EtMeNSO_2-$).

25

R^9 and R^{10} each independently represent tetrazolyl or a 5- to 6-membered heterocyclic ring comprising from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulphur, which heterocyclic ring is substituted by at least one (e.g. one, two or three) substituent selected from hydroxyl, $=O$ and $=S$.


In an embodiment of the invention R^9 and R^{10} each independently represent tetrazolyl.

30

When R^9 and R^{10} each independently represent a 5- to 6-membered heterocyclic ring, nitrogen heteroatoms in the heterocyclic ring may carry hydroxyl substituents and sulphur

atoms in the ring may be in the form of S, SO (i.e. carrying one =O substituent) or SO₂ (i.e. carrying two =O substituents).

Where R⁹ or R¹⁰ represents a 5- to 6-membered heterocyclic ring comprising from 1-4
 5 heteroatoms independently selected from nitrogen, oxygen and sulphur, which heterocyclic ring is substituted by at least one substituent selected from hydroxyl, =O and =S, examples include:



- 10 In an embodiment of the invention, M represents a bond or oxygen. In another embodiment of the invention, M represents a bond.

In an embodiment of the invention, X represents oxygen, or C₁₋₆, preferably C₁₋₄, alkylene.

- 15 In an embodiment of the invention, p, q, s and t each independently represent 2. In another embodiment of the invention, p, q, s and t each independently represent 0.


In an embodiment of the invention, where Ar¹ is a group (V) and Ar² is a thiazolyl group substituted by NH₂ and NHR⁸, R⁸ is not C₁-C₆ alkylcarbonyl or C₁-C₆ alkoxy carbonyl.

R^{28} and R^{29} together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring, which heterocyclic ring is substituted with at least one substituent (e.g. one, two or three) independently selected from CO_2R^6 , MC_{1-6} alkyl CO_2R^7 , C_{1-6} alkylsulphonylaminocarbonyl, $C(O)NHOH$, NHR^8 , R^9 and XR^{10} , and which 3- to 8-membered saturated heterocyclic ring can further be optionally substituted by at least one substituent (e.g. zero, one, two or three) independently selected from hydroxyl, halogen (e.g. fluorine, chlorine, bromine or iodine), C_1 - C_6 alkoxy (e.g. methoxy, ethoxy, n-propoxy, n-butoxy, n-pentoxo or n-hexoxy) optionally substituted by at least one halogen, and a C_1 - C_6 alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) which C_1 - C_6 alkyl group can be optionally substituted (e.g. zero, one, two or three) by at least one substituent independently selected from halogen and hydroxyl. Examples of saturated heterocyclic rings that R^{28} and R^{29} together with the nitrogen atom to which they are attached may form are rings containing one or two nitrogen atoms, e.g. pyrrolidinyl, piperidinyl, piperazinyl, homopiperazinyl, homopiperidinyl and azetidiny.

R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} and R^{25} each independently represent a hydrogen atom or a group selected from C_1 - C_6 , preferably C_1 - C_4 , alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) and C_1 - C_6 , preferably C_1 - C_4 , alkoxy (e.g. methoxy, ethoxy, n-propoxy, n-butoxy, n-pentoxo or n-hexoxy), which C_1 - C_6 alkyl or C_1 - C_6 alkoxy group can be optionally substituted with at least one (e.g. zero, one, two or three) substituent selected from halogen (e.g. fluorine, chlorine, bromine or iodine) and hydroxyl.

In a further aspect of the invention, there is provided a compound of general formula (IB), or a pharmaceutically acceptable salt or solvate thereof,

11




wherein m represents 1, 2 or 3;


each R¹ independently represents a hydrogen atom or a halogen;

A represents C(O)NH or NHC(O);


5 Ar¹ represents a group



(IIB)




(IIIB)



(IVB)

or



(VB)

10 one of R² and R³ represents halogen, nitro, NR⁴R⁵, hydroxyl, or a group selected from (i) C₁-C₆ alkyl optionally substituted by at least one halogen and (ii) C₁-C₆ alkoxy optionally substituted by at least one halogen, and the other of R² and R³ represents a hydrogen atom, halogen or a C₁-C₆ alkyl group optionally substituted by at least one halogen;

15 R⁴ and R⁵ each independently represent a hydrogen atom or a group selected from C₁-C₆ alkyl and C₁-C₆ alkoxy, which C₁-C₆ alkyl or C₁-C₆ alkoxy group can be optionally substituted with at least one substituent selected from halogen and hydroxyl;

Ar² represents phenyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, which phenyl or heteroaromatic ring is substituted by at least one substituent selected from CO₂R⁶, MC₁₋₆ alkylCO₂R⁷, C₁₋₆ alkylsulphonylaminocarbonyl, NHR⁸, R⁹ and XR¹⁰,

20 and which phenyl or heteroaromatic ring can further be optionally substituted by at least one substituent selected from halogen, nitro, NR¹¹R¹², hydroxyl, S(O)_pR¹³, a C₁-C₆ alkoxy

group which C₁-C₆ alkoxy group can be optionally substituted by at least one halogen, and a C₁-C₆ alkyl group which C₁-C₆ alkyl group can be optionally substituted by at least one substituent selected from halogen, hydroxyl, NR¹⁴R¹⁵, SO₂NR¹⁶R¹⁷, NR¹⁸SO₂R¹⁹, NHCOR²⁰ and CONR²¹R²²;

5 R⁶ and R⁷ each independently represent a hydrogen atom or a C₁-C₆ alkyl group;

R⁸ represents CN, C₁-C₆ alkylsulphonyl, C₁-C₆ alkylcarbonyl, C₁-C₆ alkoxycarbonyl, C₁-C₆ alkylaminosulphonyl, or (di)-C₁-C₆ alkylaminosulphonyl;

R⁹ and R¹⁰ each independently represent tetrazolyl or a 5- to 6-membered heterocyclic ring comprising from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and
10 sulphur, which heterocyclic ring is substituted by at least one substituent selected from hydroxyl, =O and =S;

M represents a bond, oxygen, S(O)_q or NR²³;

X represents oxygen, S(O)_s, NR²⁴, C₁-C₆ alkylene, O(CH₂)₁₋₆, NR²⁵(CH₂)₁₋₆, or S(O)_t(CH₂)₁₋₆;


15 p, q, s and t each independently represent 0, 1 or 2; and

R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴ and R²⁵ each independently represent a hydrogen atom or a group selected from C₁-C₆ alkyl and C₁-C₆ alkoxy, which C₁-C₆ alkyl or C₁-C₆ alkoxy group can be optionally substituted with at least one substituent selected from halogen and hydroxyl;

20 provided that when m is 1 and Ar¹ is a group (II) and Ar² is phenyl substituted by XR¹⁰ in a position para to Ar¹ and X is CH₂, then R¹⁰ is not a 2,4-dioxothiazolyl group; and when m is 1 and Ar¹ is a group (II) and Ar² is phenyl substituted by MC₁-C₆ alkylCO₂R⁷ in a position para to Ar¹, then M does not represent a bond.

25 In a still further aspect of the present invention there is provided a compound of formula (IC), or a pharmaceutically acceptable salt or solvate thereof,

13




wherein m represents 1, 2 or 3;

each R¹ independently represents a hydrogen atom;

A represents C(O)NH or NHC(O);

5 Ar¹ represents a group



wherein R² represents halogen, nitro, NH₂, hydroxyl or a C₁-C₆ alkyl optionally

10 substituted by one to three halogen atoms;

Ar² represents phenyl, thienyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 nitrogen atoms, wherein Ar² is substituted by at least one substituent selected from carboxyl, -C₁-C₆alkylCO₂H, -OC₁-C₆alkylCO₂H, -NHC₁-C₆alkylCO₂H, -N(C₁-C₄alkyl)C₁-C₆alkylCO₂H, -NHCN, -NHCOC₁-C₆alkyl, -NH₂SO₂C₁-C₆alkyl, -CONH₂SO₂C₁-C₆alkyl, 15 tetrazolyl, -OC₁-C₆alkyltetrazolyl and NR²⁸R²⁹, and wherein Ar² can further be optionally substituted by at least one substituent selected from halogen, trifluoromethyl, C₁-C₆alkoxy and a C₁-C₆alkyl group; and

R²⁸ and R²⁹ together with the nitrogen atom to which they are attached form a saturated heterocyclic group selected from azetidiny, pyrrolidinyl or piperidinyl, which heterocyclic 20 group is substituted by carboxyl and can further be optionally substituted by hydroxyl; provided that when Ar¹ is a group (IIC) and Ar² is phenyl substituted by C₁-C₆ alkylCO₂H in a position para to Ar¹, then m is not 1.

In an embodiment of the invention, the compound of formula (I) is selected from

4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-4-carboxylic acid,

5 4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-3-carboxylic acid,

4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

2-Chloro-5-[6-(cyanoamino)pyrazinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(cyanamino)pyrazinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

3-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyrazinecarboxylic acid,

15 3-[5-Chloro-4'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-2-pyridinyl]-benzoic acid,

2-Chloro-5-[3-[(methylsulfonyl)amino]pyrazinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(1-H-tetrazol-5-yl)pyrazinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid,

5-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid,

25 2-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-pyridinecarboxylic acid,

2-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-6-methyl-3-pyridinecarboxylic acid,

(2*S*)-2-[[[4'-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid,

30 [[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-acetic acid,

3-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid,

5-Chloro-2-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid,

5 4'-Chloro-6-methyl-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-thiophenecarboxylic acid,

6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid,

10 3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid,

2-Chloro-5-[2-(1*H*-tetrazol-5-yl)-3-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

15 2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-oxazolecarboxylic acid,

4'-Chloro-4-methyl-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-*N*-(methylsulfonyl)-2-pyridinecarboxamide,

20 *N*-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-glycine,

2-Chloro-5-[6-[(methylsulfonyl)amino]-2-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

25 [[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]oxy]-acetic acid,

2-Chloro-5-[3-(1*H*-tetrazol-5-ylmethoxy)-2-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

4'-Chloro-4-methoxy-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

30 4-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-1-methyl-1*H*-pyrazole-3-carboxylic acid,

4-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-1-methyl-1*H*-pyrazole-5-carboxylic acid,

N-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]pyrazinyl]-*N*-methyl-glycine,

5 1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]pyrazinyl]-4-piperidinecarboxylic acid,

4'-Chloro-6-fluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

4'-Chloro-5-fluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

10 4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-acetic acid,

[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-3-yl]oxy]-acetic acid,

15 (2*R*)-2-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid,

[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-4-yl]oxy]-acetic acid,

(2*S*)-2-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-3-yl]oxy]-propanoic acid,

20 4,4'-Dichloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

(2*S*)-2-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-4-yl]oxy]-propanoic acid,

25 3-Chloro-6-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid,

3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-pyridinecarboxylic acid,

30 [[2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinyl]oxy]-acetic acid,

N-[2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinyl]-glycine,

4'-Chloro-4,5-difluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

4'-Chloro-3'-[[[(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

5 3-[4-Chloro-3'-[[[(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid,

4'-Chloro-4-fluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

10 2-[5-Chloro-4'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-2-pyridinyl]-benzoic acid,

2-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-methyl-3-pyridinecarboxylic acid,

6-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-[(2-hydroxyethyl)methylamino]-3-pyridinecarboxylic acid,

15 3-[4-Methyl-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid,

4-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-1,3-dimethyl-1*H*-pyrazole-5-carboxylic acid,

20 2-[4-Methyl-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid,

3-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridineacetic acid,

1-[3-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-4-piperidinecarboxylic acid,

25 1-[3-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-L-proline,

1-[3-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-3-piperidinecarboxylic acid,

30 1-[3-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-3-azetidinecarboxylic acid,

3-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-6-methyl-2-pyridinecarboxylic acid,

3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-5-methyl-2-pyridinecarboxylic acid,

1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-4-hydroxy-4-piperidinecarboxylic acid,

5 1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-5-fluoro-2-pyridinyl]-4-piperidinecarboxylic acid,

4'-Methyl-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

10 1-[3-[4-Methyl-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-4-piperidinecarboxylic acid,

6-[4-Methyl-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid,

4-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid,

15 6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-methyl-2-pyridinecarboxylic acid,

6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-(trifluoromethyl)-2-pyridinecarboxylic acid, or


20 5-[6-(Acetylamino)-2-methyl-3-pyridinyl]-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

or a pharmaceutically acceptable salt or solvate thereof.


The present invention further provides a process for the preparation of a compound of formula (I) as defined above, or a pharmaceutically acceptable salt or solvate thereof, which comprises

25

(a) reacting a compound of formula




with a compound of formula



5 wherein one of Y and Z represents a displaceable group such as a metallic, organometallic or organosilicon group (e.g. copper, lithium, an organoboron group such as $B(OH)_2$, $B(O^iPr)_2$, BEt_2 or a boronic acid pinacol cyclic ester, or an organotin group such as $SnMe_3$ or $SnBu_3$, an organosilicon group such as $Si(Me)F_2$, an organoaluminium group such as $AlEt_2$, an organomagnesium group such as $MgCl$, $MgBr$ or MgI , or an organozinc group
 10 such as $ZnCl$, $ZnBr$ or ZnI) and the other of Y and Z represents a leaving group such as a halogeno or sulphonyloxy group (e.g. a chloro, bromo, iodo, trifluoromethanesulphonyloxy, methanesulphonyloxy or paratoluenesulphonyloxy group) and R^1 , m , A , Ar^2 , R^2 and R^3 are as defined for formula (I); or


15 (b) when Ar^2 is substituted by carboxyl, reacting a compound of formula (VI)-(IX) as defined in (a) above with a compound of formula




wherein Z is as defined in formula (X), and Ar^{2a} represents a phenyl or 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, followed by reaction with a base such as sodium hydroxide or lithium hydroxide in a solvent such as water, acetonitrile or methanol, at a temperature in the range 0-150°C, optionally followed by reaction with an acid such as hydrochloric acid in a solvent such as water, at a temperature in the range 0-150°C; or

(c) when R⁹ represents tetrazolyl, reacting a compound of formula (VI)-(IX) as defined in (a) above with a compound of formula (XI) as defined in (b) above, followed by reaction with a suitable source of azide (e.g. sodium azide, ammonium azide, azidotrimethylsilane or azidotributyltin); or

(d) when R⁸ represents CN, C₁₋₆ alkylsulphonyl, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxy carbonyl, C₁₋₆ alkylaminosulphonyl, or (di)-C₁₋₆ alkylaminosulphonyl, reacting a compound of formula (VI) – (IX) as defined in (a) above with a compound of formula



wherein L¹ represents a leaving group such as a halogeno or sulphonyloxy group (e.g. a chloro, bromo, iodo, trifluoromethanesulphonyloxy, methanesulphonyloxy or paratoluenesulphonyloxy group), Ar^{2b} represents phenyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, and Z is as defined in formula (X), followed by reaction with a compound of formula




wherein V represents a hydrogen or a metallic group, for example sodium, or

(e) when Ar^2 is substituted by carboxyl, reacting a compound of formula (VI) - (IX) as defined in (a) above with a compound of formula (XII) as defined in (d) above, followed by reaction with a suitable source of cyanide (e.g. sodium cyanide, potassium cyanide, copper cyanide or zinc cyanide), followed by reaction with a base such as sodium hydroxide or lithium hydroxide in a solvent such as water, acetonitrile or methanol, at a temperature in the range 0-150°C, optionally followed by reaction with an acid such as hydrochloric acid in a solvent such as water, at a temperature in the range 0-150°C; or

(f) when R^9 represents tetrazolyl, reacting a compound of formula (VI)-(IX) as defined in (a) above with a compound of formula (XII) as defined in (d) above, followed by reaction with a suitable source of cyanide (e.g. sodium cyanide, potassium cyanide, copper cyanide or zinc cyanide), followed by reaction with a suitable source of azide (e.g. sodium azide, ammonium azide, azidotrimethylsilane or azidotributyltin); or


(g) when Ar^2 is substituted by carboxyl, reacting a compound of formula (VI)-(IX) as defined in (a) above with a compound of formula (XII) as defined in (d) above, followed by reaction with carbon monoxide and an alcohol in the presence of a suitable catalyst, for example a palladium catalyst, followed by reaction with a base such as sodium hydroxide or lithium hydroxide in a solvent such as water or methanol, at a temperature in the range 0-150°C; or

(h) when Ar^2 represents a group of formula



(XIX),

reacting a compound of formula



with a suitable cyclodehydrating reagent (e.g. diethylaminosulfur trifluoride), followed by reaction with a suitable oxidising reagent (e.g. bromotrichloromethane and 1,8-diazabicyclo[5.4.0]undec-7-ene), followed by reaction with a base such as sodium hydroxide or lithium hydroxide in a solvent such as water or methanol, at a temperature in the range 0-150°C; or

(i) when M represents oxygen or NR^{23} , reacting a compound of formula (VI) – (IX) as defined in (a) above, with a compound of formula (XII) as defined in (d) above, followed by reaction with a compound of formula




wherein M represents oxygen or NR^{23} , and R^{23} and R^7 are as defined in formula (I), optionally followed by reaction with a base such as sodium hydroxide or lithium hydroxide in a solvent such as water or methanol, at a temperature in the range 0-150°C, or optionally followed by reaction with an acid such as hydrochloric acid, hydrobromic acid or trifluoroacetic acid in a solvent such as water, 1,4-dioxane, tetrahydrofuran, acetic acid or dichloromethane, at a temperature in the range 0-150°C; or

(j) when M represents oxygen or NR^{23} , reacting a compound of formula (XXI) as defined in (i) above, with a compound of formula (XII) as defined in (d) above, followed by

reaction with a compound of formula (VI)-(IX) as defined in (a) above, optionally followed by reaction with a base such as sodium hydroxide or lithium hydroxide in a solvent such as water or methanol, at a temperature in the range 0-150°C, or optionally followed by reaction with an acid such as hydrochloric acid, hydrobromic acid or trifluoroacetic acid in a solvent such as water, 1,4-dioxane, tetrahydrofuran, acetic acid or dichloromethane, at a temperature in the range 0-150°C; or

(k) when M represents oxygen or NR^{23} , reacting a compound of formula (VI)-(IX) as defined in (a) above, with a compound of formula




wherein Ar^{2c} represents phenyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, Z is as defined in formula (X), and M represents oxygen or NR^{23} , wherein R^{23} is as defined in formula (I), followed by reaction with either β -propiolactone or a compound of formula




wherein R^7 is as defined in formula (I), and L^1 is as defined in formula (XII), optionally followed by reaction with a base such as sodium hydroxide or lithium hydroxide in a solvent such as water or methanol, at a temperature in the range 0-150°C, or optionally followed by reaction with an acid such as hydrochloric acid, hydrobromic acid or trifluoroacetic acid in a solvent such as water, 1,4-dioxane, tetrahydrofuran, acetic acid or dichloromethane, at a temperature in the range 0-150°C; or

(l) when X represents $\text{O}(\text{CH}_2)_{1-6}$ or $\text{NR}^{25}(\text{CH}_2)_{1-6}$ and R^{10} represents tetrazolyl, reacting a compound of formula (VI)-(IX) as defined in (a) above, with a compound of formula




wherein M^1 represents oxygen or NR^{25} , R^{25} is as defined in formula (I), Ar^{2d} represents a phenyl, 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, and Z is as defined in formula (X), followed by reaction with a compound of formula




wherein L^1 is as defined in formula (XII), followed by reaction with a suitable source of azide (e.g. sodium azide, ammonium azide, azidotrimethylsilane or azidotributyltin);

(m) when Ar^2 is substituted by carboxyl, reacting a compound of formula (VI)-(IX) as defined in (a) above with a compound of formula




wherein Z is as defined in formula (X), and Ar^{2e} represents a phenyl or 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, followed by reaction with an oxidising agent such as potassium peroxymonosulfate or sodium chlorite in a solvent such as *N,N*-dimethylformamide at a temperature in the range 0-100°C; or


(n) reacting a compound of formula



25




with a compound of formula



wherein one of R^{30} and R^{31} represents NH_2 and the other of R^{30} and R^{31} represents CO_2H , $COBr$ or $COCl$, and Ar^2 , R^1 , R^2 , R^3 , R^6 and m are as defined in formula (I); or

- 10 (o) when R^{28} and R^{29} together with the nitrogen to which they are attached form a 3- to 8-membered saturated heterocyclic ring, which heterocyclic ring is substituted by CO_2R^6 , reacting a compound of formula (VI) – (IX) as defined in (a) above, with a compound of formula (XII) as defined in (d) above, followed by reaction with a compound of formula



wherein R^6 , R^{28} and R^{29} are as defined in formula (I), optionally followed by reaction with a base such as sodium hydroxide or lithium hydroxide in a solvent such as water or methanol, at a temperature in the range 0-150°C, or optionally followed by reaction with an acid such as hydrochloric acid, hydrobromic acid or trifluoroacetic acid in a solvent such

as water, 1,4-dioxane, tetrahydrofuran, acetic acid or dichloromethane, at a temperature in the range 0-150°C; or

(p) when R^{28} and R^{29} together with the nitrogen to which they are attached form a 3- to 8-membered saturated heterocyclic ring, which heterocyclic ring is substituted by CO_2R^6 , reacting a compound of formula (XII) as defined in (d) above with a compound of formula (XXXIV) as defined in (o) above, followed by reaction with a compound of formula (VI) – (IX) as defined in (a) above, optionally followed by reaction with a base such as sodium hydroxide or lithium hydroxide in a solvent such as water or methanol, at a temperature in the range 0-150°C, or optionally followed by reaction with an acid such as hydrochloric acid, hydrobromic acid or trifluoroacetic acid in a solvent such as water, 1,4-dioxane, tetrahydrofuran, acetic acid or dichloromethane, at a temperature in the range 0-150°C;

and optionally after (a), (b), (c), (d), (e), (f), (g), (h), (i), (j), (k), (l), (m), (n), (o) or (p), carrying out one or more of the following:

- converting the compound to a further compound of the invention
- forming a pharmaceutically acceptable salt or solvate of the compound.

In formula (XI), (XII), (XXII), (XXIV) and (XXVIII) above, Ar^{2a} , Ar^{2b} , Ar^{2c} , Ar^{2d} and Ar^{2e} , which independently represent phenyl or a 5- or 6-membered heteroaromatic ring, can further be optionally substituted with at least one substituent, which at least one substituent is as defined in formula (I) for further optional substituents on Ar^2 .

Where Ar^2 is substituted by CO_2R^6 , $MC_1C_6alkylCO_2R^7$ or $NR^{27}R^{28}$ wherein R^{27} and R^{28} together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring, which heterocyclic ring is substituted by CO_2R^6 or $MC_1C_6alkylCO_2R^7$, a compound of the invention wherein R^6 or R^7 represent a C_1 - C_6 alkyl group may be converted into a compound of the invention wherein R^6 or R^7 represents hydrogen by reaction with a base such as sodium hydroxide or lithium hydroxide in a solvent such as water or methanol, at a temperature in the range 0-150°C, or by reaction with an acid such as hydrochloric acid, hydrobromic acid or trifluoroacetic acid in a


solvent such as water, 1,4-dioxane, tetrahydrofuran, acetic acid or dichloromethane, at a temperature in the range 0-150°C.

Where Ar² is substituted by carboxyl, a compound of the invention may be converted into a compound of the invention where Ar² is substituted by C₁-C₆alkylsulphonylaminocarbonyl by reaction with, for example, methanesulfonamide in the presence of reagents such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 4-dimethylaminopyridine).


Compounds of formula (VI) – (IX), wherein Y represents an organoboron group such as B(OH)₂ or B(OⁱPr)₂, may be prepared by reacting compounds of formula (VI) – (IX), wherein Y represents a displaceable group such as bromo or iodo, with suitable organometallic reagents, for example methyllithium and *tert*-butyllithium, in the presence of a trialkylborate, e.g. triisopropylborate, in the presence of a suitable solvent such as tetrahydrofuran, and at a temperature in the range -100°C-30°C, and optionally followed by hydrolysis of the boronate ester by reaction with an acid such as ammonium chloride in a solvent such as water or tetrahydrofuran, at a temperature in the range 0-150 °C.

Alternatively, compounds of formula (VI) – (IX), wherein Y represents an organoboron group such as B(OH)₂ or a boronic acid pinacol cyclic ester may be prepared by reacting compounds of formula (VI) – (IX), wherein Y represents a displaceable group such as a halogeno or sulphonyloxy group, for example a chloro, bromo, iodo, trifluoromethanesulphonyloxy, methanesulphonyloxy or paratoluenesulphonyloxy group, with a suitable diboron reagent, e.g. bis(pinacolato)diboron, in the presence of a catalyst, for example palladium acetate or [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) chloride, in the presence of a base such as potassium acetate or tripotassium phosphate, in the presence of a suitable solvent, e.g. dimethylsulphoxide, 1,4-dioxane or tetrahydrofuran, and at a temperature in the range 25-250°C, and optionally followed by hydrolysis of the boronate ester by reaction with an acid such as ammonium chloride in a solvent such as water or tetrahydrofuran, at a temperature in the range 0-150 °C.


Compounds of formula (VI) – (IX), wherein Y represents a leaving group such as a halogeno or sulphonyloxy group, may conveniently be prepared by reacting a compound of general formula



5 with a compound of general formula




10 wherein one of R^{26} and R^{27} represents NH_2 and the other of R^{26} and R^{27} represents CO_2H , $COBr$ or $COCl$, R^1 and m are as defined in formula (I), Y represents a leaving group such as a halogeno or sulphonyloxy group as defined in formulae (VI) – (IX), and R^2 and R^3 are as defined in formula (I), optionally in the presence of suitable coupling reagents such as 1,1'-carbonyldiimidazole or dicyclohexylcarbodiimide and 1-hydroxybenzotriazole. Compounds of formula (XX) may be prepared by reacting a compound of formula



with L-serine methyl ester, in the presence of suitable coupling reagents such as 1,1'-carbonyldiimidazole or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 1-hydroxybenzotriazole.





- 5 Compounds of formula (XXVI) may be prepared by reacting a compound of formula



with a suitable oxidant such as potassium peroxomonosulfate or sodium chlorite.

10


Compounds of formula (XXIX) – (XXXII) where R³⁰ is a carboxyl group may be prepared by reacting a compound of general formula



15

wherein Ar^2 , R^2 , and R^3 are as defined in formula (I), with an acid such as hydrochloric acid, hydrobromic acid or trifluoroacetic acid in a solvent such as water, 1,4-dioxane, tetrahydrofuran, acetic acid or dichloromethane, at a temperature in the range 0-150°C.

- 5 Compounds of formula (XXXV) – (XXXVIII) may be prepared by reacting a compound of formula



10


- with a compound of formula (X) as defined in (a) above wherein Y is as defined in formula (VI)-(IX), and R^2 and R^3 are as defined in formula (I), in the presence of a catalyst such as tetrakis(triphenylphosphine)palladium(0), palladium(II) chloride, palladium(II) bromide, dichlorobis(triphenylphosphine)palladium(II), nickel(II) chloride, nickel(II) bromide or
- 15 bis(triphenylphosphine)nickel(II) chloride, in the presence of a suitable solvent such as tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane, benzene, toluene, xylene, methanol, ethanol or water. The reaction may be conducted in the presence of a suitable base such as sodium carbonate or potassium carbonate, pyridine, 4-dimethylaminopyridine, triethylamine or morpholine, and at a temperature in the range 10 to 250°C, e.g. in the
- 20 range 60 to 120°C.


Compounds of formula (XXXIX) – (XXXXXII), wherein Y represents an organoboron group such as $B(OH)_2$ or $B(O^iPr)_2$, may be prepared by reacting compounds of formula (XXXIX) – (XXXXXII), wherein Y represents a displaceable group such as bromo or iodo,

with suitable organometallic reagents, for example methyllithium and *tert*-butyllithium, in the presence of a trialkylborate, e.g. triisopropylborate, in the presence of a suitable solvent such as tetrahydrofuran, and at a temperature in the range -100°C to 30°C, and optionally followed by hydrolysis of the boronate ester by reaction with an acid such as ammonium chloride in a solvent such as water or tetrahydrofuran, at a temperature in the range 0-150 °C.

Alternatively, compounds of formula (XXXIX) – (XXXXII), wherein Y represents an organoboron group such as B(OH)₂ or a boronic acid pinacol cyclic ester may be prepared by reacting compounds of formula (XXXIX) – (XXXXII), wherein Y represents a displaceable group such as a halogeno or sulphonyloxy group, for example a chloro, bromo, iodo, trifluoromethanesulphonyloxy, methanesulphonyloxy or paratoluenesulphonyloxy group, with a suitable diboron reagent, e.g. bis(pinacolato)diboron, in the presence of a catalyst, for example palladium acetate or [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) chloride, in the presence of a base such as potassium acetate or tripotassium phosphate, in the presence of a suitable solvent, e.g. dimethylsulphoxide, 1,4-dioxane or tetrahydrofuran, and at a temperature in the range 25-250°C, and optionally followed by hydrolysis of the boronate ester by reaction with an acid such as ammonium chloride in a solvent such as water or tetrahydrofuran, at a temperature in the range 0-150 °C.

Compounds of formula (XXXIX) – (XXXXII), wherein Y represents a leaving group such as a halogeno or sulphonyloxy group, may conveniently be prepared by reacting a compound of formula





wherein R³² represents CO₂H, COBr or COCl, Y is a leaving group as defined in formula (VI) – (IX), and R² and R³ are as defined in formula (I), with *tert*-butanol or potassium *tert*-butoxide, optionally in the presence of suitable reagents such as dicyclohexylcarbodiimide and 4-dimethylaminopyridine.

Compounds of formula (X), (XI), (XII), (XIII), (XIV), (XV), (XVI), (XVII), (XVIII), (XXI), (XXII), (XXIII), (XXIV), (XXV), (XXVII), (XXVIII), (XXXIII), (XXXIV), (XXXXIII), (XXXXIV), (XXXXXV) and (XXXXXVI) are either commercially available, are known in the literature or may be prepared easily using known techniques.

In processes (a), (b), (c), (d), (e), (f), (g), (i), (j), (k), (l), (m), (o) and (p), the coupling reaction is conveniently carried out in the presence of a catalyst such as tetrakis(triphenylphosphine)palladium(0), palladium(II) chloride, palladium(II) bromide, dichlorobis(triphenylphosphine)palladium(II), nickel(II) chloride, nickel(II) bromide or bis(triphenylphosphine)nickel(II) chloride, in the presence of a suitable solvent such as tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane, benzene, toluene, xylene, methanol, ethanol or water. The reaction is preferably conducted in the presence of a suitable base such as sodium carbonate or potassium carbonate, pyridine, 4-dimethylaminopyridine, triethylamine or morpholine, and at a temperature in the range 10 to 250°C, preferably in the range 60 to 120°C.

In processes (c), (f) and (l), the tetrazole formation reaction is carried out in the presence of a suitable source of azide, for example sodium azide, ammonium azide, azidotrimethylsilane or azidotributyltin, optionally in the presence of a suitable catalyst, for example dibutyltin oxide, in the presence of a suitable solvent such as toluene, *N,N*-

dimethylformamide or 1-methyl-2-pyrrolidinone, and at a temperature in the range 10 to 250°C, preferably in the range 50 to 120°C.

In process (d), the displacement reaction may be carried out in the presence of a suitable
5 base, for example potassium *tert*-butoxide, sodium hydride, potassium carbonate or caesium carbonate, optionally in the presence of a suitable catalyst, for example a palladium catalyst such as tetrakis(triphenylphosphine)palladium(0), palladium(II) chloride, palladium(II) bromide, palladium(II) acetate,
dichlorobis(triphenylphosphine)palladium(II) or tris(dibenzylideneacetone)palladium(0),
10 or a copper catalyst such as copper(I) iodide, optionally in the presence of a suitable ligand, for example 1,1'-bis(diphenylphosphino)ferrocene, 9,9-dimethyl-4,5-bis(diphenylphosphino)xanthene or 2-dicyclohexylphosphino-2'-(*N,N*-dimethylamino)biphenyl, in the presence of a suitable solvent, for example 1-methyl-2-pyrrolidinone, 1,4-dioxane, 1,2-dimethoxyethane, tetrahydrofuran or acetonitrile, and at a
15 temperature in the range 10 to 250°C, preferably in the range 60 to 150°C.

In processes (e) and (f), the displacement reaction may be carried out in the presence of a suitable source of cyanide, for example sodium cyanide, potassium cyanide, copper cyanide or zinc cyanide, optionally in the presence of a suitable catalyst, for example a
20 palladium catalyst such as tetrakis(triphenylphosphine)palladium(0) or palladium(II) acetate, in the presence of a suitable solvent, for example *N,N*-dimethylformamide, 1-methyl-2-pyrrolidinone or dimethylsulfoxide, and at a temperature in the range 10-250°C, preferably in the range 60 to 150°C.

25 In process (g), the carbonylation reaction may be carried out in the presence of an alcohol such as butanol, propanol, ethanol or methanol, in the presence of a catalyst such as tetrakis(triphenylphosphine)palladium(0), palladium(II) chloride, palladium(II) bromide, palladium(II) acetate, dichlorobis(triphenylphosphine)palladium (II) or [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) chloride, optionally in the presence of a
30 ligand such as triphenylphosphine or 1,3-bis(diphenylphosphino)propane, in the presence of a suitable base, for example triethylamine, optionally in the presence of a co-solvent, for example 1-methyl-2-pyrrolidinone or *N,N*-dimethylformamide, and at a temperature in the

range 10-150°C, and under a pressure of carbon monoxide in the range of 100 kPa to 1000 kPa (1 to 10bar).

In process (h), the cyclodehydration reaction may be carried out in the presence of a cyclodehydrating reagent such as diethylaminosulfur trifluoride or Burgess reagent ((methoxycarbonylsulfamoyl)triethylammonium hydroxide, inner salt), in the presence of a suitable solvent, for example dichloromethane, and at a temperature in the range -78-30°C. The oxidation reaction may be carried out in the presence of oxidising reagents such as bromotrichloromethane and 1,8-diazabicyclo[5.4.0]undec-7-ene, or manganese dioxide, in the presence of a suitable solvent such as dichloromethane, at a temperature in the range 0-150°C.

In processes (i) and (j), the displacement reaction may optionally be carried out in the presence of a suitable catalyst such as palladium(II) acetate, optionally in the presence of a suitable ligand such as 9,9-dimethyl-4,5-bis(diphenylphosphino)xanthene or 2-dicyclohexylphosphino-2'-(*N,N*-dimethylamino)biphenyl, in the presence of a suitable base, for example potassium *tert*-butoxide, sodium *tert*-butoxide, triethylamine or potassium carbonate, in the presence of a suitable solvent, for example tetrahydrofuran, acetonitrile, *N*-methylpyrrolidinone, toluene or acetone, and at a temperature in the range 0-150°C.

In processes (k) and (l), the displacement reaction may be carried out in the presence of a suitable base, for example potassium *tert*-butoxide, triethylamine or potassium carbonate, in the presence of a suitable solvent, for example tetrahydrofuran, acetonitrile, *N*-methylpyrrolidinone or acetone, and at a temperature in the range 0-150°C.

In process (m), the oxidation reaction may be carried out in the presence of an oxidising agent such as potassium peroxymonosulfate or sodium chlorite in a solvent such as *N,N*-dimethylformamide at a temperature in the range 0-100°C.

In process (n), the amide coupling reaction may be carried out in the presence of a suitable coupling reagent, such as 1,1'-carbonyldiimidazole or dicyclohexylcarbodiimide and 1-hydroxybenzotriazole, in the presence of a base such as triethylamine, *N*-methylmorpholine, diisopropylethylamine or potassium carbonate, in a solvent such as dichloromethane, *N*-methylpyrrolidinone, *N,N*-dimethylformamide or tetrahydrofuran, and at a temperature in the range 0-150°C.

In processes (o) and (p), the displacement reaction may optionally be carried out in the presence of a suitable base, such as triethylamine, in a solvent such as acetonitrile or pyridine, and at a temperature in the range 0-150°C.

It will be appreciated by those skilled in the art that in the processes of the present invention certain functional groups such as hydroxyl, carboxyl or amino groups in the starting reagents or intermediate compounds may need to be protected by protecting groups. Thus, the preparation of the compounds of formula (I) may involve at a certain stage the removal of one or more protecting groups.

The protection and deprotection of functional groups is described in 'Protective Groups in Organic Synthesis', 2nd edition, T.W. Greene and P.G.M. Wuts, Wiley-Interscience (1991) and 'Protecting Groups', P.J. Kocienski, Georg Thieme Verlag (1994).

The compounds of formula (I) above may be converted to a pharmaceutically acceptable salt or solvate thereof. Where the compound is sufficiently acidic, suitable salts include base salts such as an alkali metal salt for example sodium or potassium, an alkaline earth metal salt for example calcium or magnesium, an organic amine salt for example triethylamine, morpholine, *N*-methylpiperidine, *N*-ethylpiperidine, procaine, dibenzylamine, *N,N*-dibenzylethylamine or amino acids for example lysine. Where the compound is sufficiently basic, suitable salts include acid addition salts such as a hydrochloride, hydrobromide, phosphate, acetate, fumarate, maleate, tartrate, citrate, oxalate, methanesulphonate or *p*-toluenesulphonate salt. There may be more than one cation or anion depending on the number of charged functions and the valency of the

cations or anions. Other pharmaceutically acceptable salts, as well as prodrugs such as pharmaceutically acceptable esters and pharmaceutically acceptable amides may be prepared using conventional methods. It will be understood that certain compounds of the present invention may exist in solvated, for example hydrated, as well as unsolvated forms.

5 It is to be understood that the present invention encompasses all such solvated forms.

A compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, can be used in the treatment of:

1. *respiratory tract*: obstructive diseases of the airways including: asthma, including
10 bronchial, allergic, intrinsic, extrinsic, exercise-induced, drug-induced (including aspirin and NSAID-induced) and dust-induced asthma, both intermittent and persistent and of all severities, and other causes of airway hyper-responsiveness; chronic obstructive pulmonary disease (COPD); bronchitis, including infectious and eosinophilic bronchitis; emphysema; bronchiectasis; cystic fibrosis; sarcoidosis; farmer's lung and related diseases;
15 hypersensitivity pneumonitis; lung fibrosis, including cryptogenic fibrosing alveolitis, idiopathic interstitial pneumonias, fibrosis complicating anti-neoplastic therapy and chronic infection, including tuberculosis and aspergillosis and other fungal infections; complications of lung transplantation; vasculitic and thrombotic disorders of the lung vasculature, and pulmonary hypertension; antitussive activity including treatment of
20 chronic cough associated with inflammatory and secretory conditions of the airways, and iatrogenic cough; acute and chronic rhinitis including rhinitis medicamentosa, and vasomotor rhinitis; perennial and seasonal allergic rhinitis including rhinitis nervosa (hay fever); nasal polyposis; acute viral infection including the common cold, and infection due to respiratory syncytial virus, influenza, coronavirus (including SARS) and adenovirus;
- 25 2. *bone and joints*: arthritides associated with or including osteoarthritis/osteoarthrosis, both primary and secondary to, for example, congenital hip dysplasia; cervical and lumbar spondylitis, and low back and neck pain; rheumatoid arthritis and Still's disease; seronegative spondyloarthropathies including ankylosing spondylitis, psoriatic arthritis, reactive arthritis and undifferentiated spondarthropathy; septic arthritis and other infection-
30 related arthropathies and bone disorders such as tuberculosis, including Potts' disease and Poncet's syndrome; acute and chronic crystal-induced synovitis including urate gout, calcium pyrophosphate deposition disease, and calcium apatite related tendon, bursal and

synovial inflammation; Behcet's disease; primary and secondary Sjogren's syndrome; systemic sclerosis and limited scleroderma; systemic lupus erythematosus, mixed connective tissue disease, and undifferentiated connective tissue disease; inflammatory myopathies including dermatomyositis and polymyositis; polymyalgia rheumatica; juvenile arthritis including idiopathic inflammatory arthritides of whatever joint distribution and associated syndromes, and rheumatic fever and its systemic complications; vasculitides including giant cell arteritis, Takayasu's arteritis, Churg-Strauss syndrome, polyarteritis nodosa, microscopic polyarteritis, and vasculitides associated with viral infection, hypersensitivity reactions, cryoglobulins, and paraproteins; low back pain; Familial Mediterranean fever, Muckle-Wells syndrome, and Familial Hibernian Fever, Kikuchi disease; drug-induced arthralgias, tendonitides, and myopathies;

3. *pain and connective tissue remodelling of musculoskeletal disorders due to injury [for example sports injury] or disease:* arthritides (for example rheumatoid arthritis, osteoarthritis, gout or crystal arthropathy), other joint disease (such as intervertebral disc degeneration or temporomandibular joint degeneration), bone remodelling disease (such as osteoporosis, Paget's disease or osteonecrosis), chondroititis, scleroderma, mixed connective tissue disorder, spondyloarthropathies or periodontal disease (such as periodontitis);

4. *skin:* psoriasis, atopic dermatitis, contact dermatitis or other eczematous dermatoses, and delayed-type hypersensitivity reactions; phyto- and photodermatitis; seborrhoeic dermatitis, dermatitis herpetiformis, lichen planus, lichen sclerosus et atrophica, pyoderma gangrenosum, skin sarcoid, discoid lupus erythematosus, pemphigus, pemphigoid, epidermolysis bullosa, urticaria, angioedema, vasculitides, toxic erythemas, cutaneous eosinophilias, alopecia areata, male-pattern baldness, Sweet's syndrome, Weber-Christian syndrome, erythema multiforme; cellulitis, both infective and non-infective; panniculitis; cutaneous lymphomas, non-melanoma skin cancer and other dysplastic lesions; drug-induced disorders including fixed drug eruptions;

5. *eyes:* blepharitis; conjunctivitis, including perennial and vernal allergic conjunctivitis; iritis; anterior and posterior uveitis; choroiditis; autoimmune; degenerative or inflammatory disorders affecting the retina; ophthalmitis including sympathetic ophthalmitis; sarcoidosis; infections including viral, fungal, and bacterial;

6. *gastrointestinal tract*: glossitis, gingivitis, periodontitis; oesophagitis, including reflux; eosinophilic gastro-enteritis, mastocytosis, Crohn's disease, colitis including ulcerative colitis, proctitis, pruritis ani; coeliac disease, irritable bowel syndrome, and food-related allergies which may have effects remote from the gut (for example migraine, rhinitis or eczema);

7. *abdominal*: hepatitis, including autoimmune, alcoholic and viral; fibrosis and cirrhosis of the liver; cholecystitis; pancreatitis, both acute and chronic;

8. *genitourinary*: nephritis including interstitial and glomerulonephritis; nephrotic syndrome; cystitis including acute and chronic (interstitial) cystitis and Hunner's ulcer; acute and chronic urethritis, prostatitis, epididymitis, oophoritis and salpingitis; vulvo-vaginitis; Peyronie's disease; erectile dysfunction (both male and female);

9. *allograft rejection*: acute and chronic following, for example, transplantation of kidney, heart, liver, lung, bone marrow, skin or cornea or following blood transfusion; or chronic graft versus host disease;

10. *CNS*: Alzheimer's disease and other dementing disorders including CJD and nvCJD; amyloidosis; multiple sclerosis and other demyelinating syndromes; cerebral atherosclerosis and vasculitis; temporal arteritis; myasthenia gravis; acute and chronic pain (acute, intermittent or persistent, whether of central or peripheral origin) including visceral pain, headache, migraine, trigeminal neuralgia, atypical facial pain, joint and bone pain, pain arising from cancer and tumor invasion, neuropathic pain syndromes including diabetic, post-herpetic, and HIV-associated neuropathies; neurosarcoidosis; central and peripheral nervous system complications of malignant, infectious or autoimmune processes;

11. *other auto-immune and allergic disorders* including Hashimoto's thyroiditis, Graves' disease, Addison's disease, diabetes mellitus, idiopathic thrombocytopaenic purpura, eosinophilic fasciitis, hyper-IgE syndrome, antiphospholipid syndrome;

12. *other disorders with an inflammatory or immunological component*; including acquired immune deficiency syndrome (AIDS), leprosy, Sezary syndrome, and paraneoplastic syndromes;

13. *cardiovascular*: atherosclerosis, affecting the coronary and peripheral circulation; pericarditis; myocarditis, inflammatory and auto-immune cardiomyopathies including myocardial sarcoid; ischaemic reperfusion injuries; endocarditis, valvulitis, and aortitis

including infective (for example syphilitic); vasculitides; disorders of the proximal and peripheral veins including phlebitis and thrombosis, including deep vein thrombosis and complications of varicose veins;

14. *oncology*: treatment of common cancers including prostate, breast, lung, ovarian, pancreatic, bowel and colon, stomach, skin and brain tumors and malignancies affecting the bone marrow (including the leukaemias) and lymphoproliferative systems, such as Hodgkin's and non-Hodgkin's lymphoma; including the prevention and treatment of metastatic disease and tumour recurrences, and paraneoplastic syndromes; and,

15. *gastrointestinal tract*: Coeliac disease, proctitis, eosinophilic gastro-enteritis, mastocytosis, Crohn's disease, ulcerative colitis, microscopic colitis, indeterminant colitis, irritable bowel disorder, irritable bowel syndrome, non-inflammatory diarrhea, food-related allergies which have effects remote from the gut, e.g., migraine, rhinitis and eczema.

Accordingly, the present invention provides a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined for use in therapy.


In another aspect, the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for use in therapy.

In the context of the present specification, the term "therapy" also includes "prophylaxis" unless there are specific indications to the contrary. The terms "therapeutic" and "therapeutically" should be construed accordingly.

The invention further provides a method of effecting immunosuppression (e.g. in the treatment of rheumatoid arthritis, osteoarthritis, irritable bowel disease, atherosclerosis or psoriasis) which comprises administering a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined to a patient.

The invention also provides a method of treating an obstructive airways disease (e.g. asthma or COPD) which comprises administering to a patient a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined to a patient.

The compounds of the present invention are especially advantageous as pharmaceuticals for use in the treatment of inflammatory disorders such as rheumatoid arthritis, osteoarthritis, asthma and chronic obstructive pulmonary disease (COPD). Accordingly, the present invention provides for the use of a compound of formula (IA), or a pharmaceutically acceptable salt or solvate thereof, in the manufacture of a medicament for use in the treatment of an inflammatory disorder




wherein m represents 1, 2 or 3;


each R¹ independently represents a hydrogen atom or a halogen;

A represents C(O)NH or NHC(O);


Ar¹ represents a group



(II)




(III)



(IV)

or



(V)

one of R^2 and R^3 represents halogen, nitro, NR^4R^5 , hydroxyl, or a group selected from (i) C_1 - C_6 alkyl optionally substituted by at least one halogen and (ii) C_1 - C_6 alkoxy optionally substituted by at least one halogen, and the other of R^2 and R^3 represents a hydrogen atom, halogen or a C_1 - C_6 alkyl group optionally substituted by at least one halogen;

R^4 and R^5 each independently represent a hydrogen atom or a group selected from C_1 - C_6 alkyl and C_1 - C_6 alkoxy, which C_1 - C_6 alkyl or C_1 - C_6 alkoxy group can be optionally substituted with at least one substituent selected from halogen and hydroxyl;

Ar^2 represents phenyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, which phenyl or heteroaromatic ring is substituted by at least one substituent selected from CO_2R^6 , $MC_{1-6}alkylCO_2R^7$, $C_{1-6}alkylsulphonylaminocarbonyl$, NHR^8 , R^9 , XR^{10} , $C(O)NHOH$ and $NR^{28}R^{29}$;

and which phenyl or heteroaromatic ring can further be optionally substituted by at least one substituent selected from halogen, nitro, $NR^{11}R^{12}$, hydroxyl, $S(O)_pR^{13}$, a C_1 - C_6 alkoxy group which C_1 - C_6 alkoxy group can be optionally substituted by a halogen, and a C_1 - C_6 alkyl group which C_1 - C_6 alkyl group can be optionally substituted by at least one substituent selected from halogen, hydroxyl, $NR^{14}R^{15}$, $SO_2NR^{16}R^{17}$, $NR^{18}SO_2R^{19}$, $NHCOR^{20}$ and $CONR^{21}R^{22}$;

R^6 and R^7 each independently represent a hydrogen atom or a C_1 - C_6 alkyl group;

R^8 represents CN , C_1 - C_6 alkylsulphonyl, C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkoxycarbonyl, C_1 - C_6 alkylaminosulphonyl, or (di)- C_1 - C_6 alkylaminosulphonyl;

R^9 and R^{10} each independently represent tetrazolyl or a 5- to 6-membered heterocyclic ring comprising from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulphur, which heterocyclic ring is substituted by at least one substituent selected from hydroxyl, $=O$ and $=S$;

M represents a bond, oxygen, $S(O)_q$ or NR^{23} ;

X represents oxygen, $S(O)_s$, NR^{24} , C_1 - C_6 alkylene, $O(CH_2)_{1-6}$, $NR^{25}(CH_2)_{1-6}$, or $S(O)_t(CH_2)_{1-6}$;

p , q , s and t each independently represent 0, 1 or 2;

R^{28} and R^{29} together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring, which heterocyclic ring is substituted with at least

one substituent independently selected from CO_2R^6 , $\text{MC}_1\text{-C}_6$ alkyl CO_2R^7 , $\text{C}_1\text{-C}_6$ alkylsulphonylaminocarbonyl, $\text{C}(\text{O})\text{NHOH}$, NHR^8 , R^9 and XR^{10} , and which 3- to 8-membered saturated heterocyclic ring can further be optionally substituted by at least one substituent independently selected from hydroxyl, halogen, $\text{C}_1\text{-C}_6$ alkoxy optionally substituted by at least one halogen, and a $\text{C}_1\text{-C}_6$ alkyl group which $\text{C}_1\text{-C}_6$ alkyl group can be optionally substituted by at least one substituent independently selected from halogen and hydroxyl; and

R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} and R^{25} each independently represent a hydrogen atom or a group selected from $\text{C}_1\text{-C}_6$ alkyl and $\text{C}_1\text{-C}_6$ alkoxy, which $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_1\text{-C}_6$ alkoxy group can be optionally substituted with at least one substituent selected from halogen and hydroxyl.

The present invention further provides for the use of compound of formula (IA), or a pharmaceutically acceptable salt or solvate thereof, in the manufacture of a medicament for use in the treatment of atherosclerosis.

The present invention further provides a method of treating an inflammatory disorder (e.g. rheumatoid arthritis, osteoarthritis, asthma or chronic obstructive pulmonary disease) or atherosclerosis, which comprises administering a therapeutically effective amount of a compound of formula (IA), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined to a patient.

For all the above-mentioned therapeutic uses the dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the disorder indicated. The daily dosage of the compound of formula (I)/salt/solvate ("active ingredient") may be in the range from 0.001 mg/kg to 30 mg/kg.

The compounds of formula (I) and pharmaceutically acceptable salts or solvates thereof may be used on their own but will generally be administered in the form of a pharmaceutical composition in which the formula (I) compound/salt/solvate ("active ingredient") is in association with a pharmaceutically acceptable adjuvant, diluent or carrier. Depending on the mode of administration, the pharmaceutical composition will

preferably comprise from 0.05 to 99 %w (per cent by weight), more preferably from 0.10 to 70 %w, of active ingredient, and, from 1 to 99.95 %w, more preferably from 30 to 99.90 %w, of a pharmaceutically acceptable adjuvant, diluent or carrier, all percentages by weight being based on total composition.

5

Thus, the present invention also provides a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

10

The pharmaceutical composition of the invention may be administered topically (e.g. to the lung and/or airways or to the skin) in the form of solutions, suspensions, heptafluoroalkane aerosols and dry powder formulations; or systemically, e.g. by oral administration in the form of tablets, capsules, syrups, powders or granules, or by parenteral administration in the form of solutions or suspensions, or by subcutaneous administration or by rectal administration in the form of suppositories or transdermally.

15

The invention further relates to combination therapies wherein a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition or formulation comprising a compound of the invention, is administered concurrently or sequentially or as a combined preparation with another therapeutic agent or agents, for the treatment of one or more of the conditions listed.

20

In particular, for the treatment of the inflammatory diseases such as (but not restricted to) rheumatoid arthritis, osteoarthritis, asthma, allergic rhinitis, chronic obstructive pulmonary disease (COPD), psoriasis, and inflammatory bowel disease, the compounds of the invention may be combined with the following agents: Non-steroidal anti-inflammatory agents (hereinafter NSAIDs) including non-selective cyclo-oxygenase COX-1 / COX-2 inhibitors whether applied topically or systemically (such as piroxicam, diclofenac, propionic acids such as naproxen, flurbiprofen, fenoprofen, ketoprofen and ibuprofen, fenamates such as mefenamic acid, indomethacin, sulindac, azapropazone, pyrazolones

30

such as phenylbutazone, salicylates such as aspirin); selective COX-2 inhibitors (such as meloxicam, celecoxib, rofecoxib, valdecoxib, lumaprocoxib, parecoxib and etoricoxib); cyclo-oxygenase inhibiting nitric oxide donors (CINODs); glucocorticosteroids (whether administered by topical, oral, intramuscular, intravenous, or intra-articular routes);
5 methotrexate; leflunomide; hydroxychloroquine; D-penicillamine; auranofin or other parenteral or oral gold preparations; analgesics; diacerein; intra-articular therapies such as hyaluronic acid derivatives; and nutritional supplements such as glucosamine.

The present invention still further relates to the combination of a compound of the
10 invention, or a pharmaceutically acceptable salt or solvate thereof, together with a cytokine or agonist or antagonist of cytokine function, (including agents which act on cytokine signalling pathways such as modulators of the SOCS system) including alpha-, beta-, and gamma-interferons; insulin-like growth factor type I (IGF-1); interleukins (IL) including IL1 to 17, and interleukin antagonists or inhibitors such as anakinra; tumour necrosis factor
15 alpha (TNF- α) inhibitors such as anti-TNF monoclonal antibodies (for example infliximab; adalimumab, and CDP-870) and TNF receptor antagonists including immunoglobulin molecules (such as etanercept) and low-molecular-weight agents such as pentoxifylline.

In addition the invention relates to a combination of a compound of the invention, or a
20 pharmaceutically acceptable salt or solvate thereof, with a monoclonal antibody targeting B-Lymphocytes (such as CD20 (rituximab), MRA- α IL16R and T-Lymphocytes, CTLA4-Ig, HuMax II-15).

The present invention still further relates to the combination of a compound of the
25 invention, or a pharmaceutically acceptable salt thereof, with a modulator of chemokine receptor function such as an antagonist of CCR1, CCR2, CCR2A, CCR2B, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CCR10 and CCR11 (for the C-C family); CXCR1, CXCR2, CXCR3, CXCR4 and CXCR5 (for the C-X-C family) and CX₃CR1 for the C-X₃-C family.

30 The present invention further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, with an inhibitor of matrix

metalloprotease (MMPs), i.e., the stromelysins, the collagenases, and the gelatinases, as well as aggrecanase; especially collagenase-1 (MMP-1), collagenase-2 (MMP-8), collagenase-3 (MMP-13), stromelysin-1 (MMP-3), stromelysin-2 (MMP-10), and stromelysin-3 (MMP-11) and MMP-9 and MMP-12, including agents such as doxycycline.

5

The present invention still further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt thereof, and a leukotriene biosynthesis inhibitor, 5-lipoxygenase (5-LO) inhibitor or 5-lipoxygenase activating protein (FLAP) antagonist such as; zileuton; ABT-761; fenleuton; tepoxalin; Abbott-79175; Abbott-85761; a N-(5-substituted)-thiophene-2-alkylsulfonamide; 2,6-di-tert-butylphenolhydrazones; a methoxytetrahydropyrans such as Zeneca ZD-2138; the compound SB-210661; a pyridinyl-substituted 2-cyanonaphthalene compound such as L-739,010; a 2-cyanoquinoline compound such as L-746,530; or an indole or quinoline compound such as MK-591, MK-886, and BAY x 1005.

15

The present invention further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and a receptor antagonist for leukotrienes (LT) B₄, LTC₄, LTD₄, and LTE₄. selected from the group consisting of the phenothiazin-3-1s such as L-651,392; amidino compounds such as CGS-25019c; benzoxalamines such as ontazolast; benzenecarboximidamides such as BIIL 284/260; and compounds such as zafirlukast, ablukast, montelukast, pranlukast, verlukast (MK-679), RG-12525, Ro-245913, iralukast (CGP 45715A), and BAY x 7195.

20

The present invention still further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and a phosphodiesterase (PDE) inhibitor such as a methylxanthanine including theophylline and aminophylline; a selective PDE isoenzyme inhibitor including a PDE4 inhibitor an inhibitor of the isoform PDE4D, or an inhibitor of PDE5.

25

The present invention further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and a histamine type 1 receptor antagonist such as cetirizine, loratadine, desloratadine, fexofenadine, acrivastine,

30

terfenadine, astemizole, azelastine, levocabastine, chlorpheniramine, promethazine, cyclizine, or mizolastine; applied orally, topically or parenterally.

The present invention still further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and a proton pump inhibitor (such as omeprazole) or a gastroprotective histamine type 2 receptor antagonist.

The present invention further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and an antagonist of the histamine type 4 receptor.

The present invention still further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and an alpha-1/alpha-2 adrenoceptor agonist vasoconstrictor sympathomimetic agent, such as propylhexedrine, phenylephrine, phenylpropanolamine, ephedrine, pseudoephedrine, naphazoline hydrochloride, oxymetazoline hydrochloride, tetrahydrozoline hydrochloride, xylometazoline hydrochloride, tramazoline hydrochloride or ethylnorepinephrine hydrochloride.

The present invention further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and an anticholinergic agents including muscarinic receptor (M1, M2, and M3) antagonist such as atropine, hyoscine, glycopyrrrolate, ipratropium bromide, tiotropium bromide, oxitropium bromide, pirenzepine or telenzepine.

The present invention still further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and a beta-adrenoceptor agonist (including beta receptor subtypes 1-4) such as isoprenaline, salbutamol, formoterol, salmeterol, terbutaline, orciprenaline, bitolterol mesylate, or pirbuterol, or a chiral enantiomer thereof.

The present invention further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and a chromone, such as sodium cromoglycate or nedocromil sodium.

5 The present invention still further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, with a glucocorticoid, such as flunisolide, triamcinolone acetonide, beclomethasone dipropionate, budesonide, fluticasone propionate, ciclesonide or mometasone furoate.

10 The present invention further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, with an agent that modulates a nuclear hormone receptor such as PPARs.

The present invention still further relates to the combination of a compound of the
15 invention, or a pharmaceutically acceptable salt or solvate thereof, together with an immunoglobulin (Ig) or Ig preparation or an antagonist or antibody modulating Ig function such as anti-IgE (for example omalizumab).

The present invention further relates to the combination of a compound of the invention, or
20 a pharmaceutically acceptable salt or solvate thereof, and another systemic or topically-applied anti-inflammatory agent, such as thalidomide or a derivative thereof, a retinoid, dithranol or calcipotriol.

The present invention still further relates to the combination of a compound of the
25 invention, or a pharmaceutically acceptable salt or solvate thereof, and combinations of aminosalicylates and sulfapyridine such as sulfasalazine, mesalazine, balsalazide, and olsalazine; and immunomodulatory agents such as the thiopurines, and corticosteroids such as budesonide.

30 The present invention further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, together with an antibacterial agent such as a penicillin derivative, a tetracycline, a macrolide, a beta-lactam, a

fluoroquinolone, metronidazole, an inhaled aminoglycoside; an antiviral agent including acyclovir, famciclovir, valaciclovir, ganciclovir, cidofovir, amantadine, rimantadine, ribavirin, zanamavir and oseltamavir; a protease inhibitor such as indinavir, nelfinavir, ritonavir, and saquinavir; a nucleoside reverse transcriptase inhibitor such as didanosine, lamivudine, stavudine, zalcitabine or zidovudine; or a non-nucleoside reverse transcriptase inhibitor such as nevirapine or efavirenz.

The present invention still further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and a cardiovascular agent such as a calcium channel blocker, a beta-adrenoceptor blocker, an angiotensin-converting enzyme (ACE) inhibitor, an angiotensin-2 receptor antagonist; a lipid lowering agent such as a statin or a fibrate; a modulator of blood cell morphology such as pentoxifylline; thrombolytic, or an anticoagulant such as a platelet aggregation inhibitor.

The present invention further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and a CNS agent such as an antidepressant (such as sertraline), an anti-Parkinsonian drug (such as deprenyl, L-dopa, ropinirole, pramipexole, a MAOB inhibitor such as selegine and rasagiline, a COMT inhibitor such as tasmar, an A-2 inhibitor, a dopamine reuptake inhibitor, an NMDA antagonist, a nicotine agonist, a dopamine agonist or an inhibitor of neuronal nitric oxide synthase), or an anti-Alzheimer's drug such as donepezil, rivastigmine, tacrine, a COX-2 inhibitor, propentofylline or metrifonate.

The present invention still further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and an agent for the treatment of acute or chronic pain, such as a centrally or peripherally-acting analgesic (for example an opioid or derivative thereof), carbamazepine, phenytoin, sodium valproate, amitriptyline or other anti-depressant agent-s, paracetamol, or a non-steroidal anti-inflammatory agent.

The present invention further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, together with a parenterally or

topically-applied (including inhaled) local anaesthetic agent such as lignocaine or a derivative thereof.

A compound of the present invention, or a pharmaceutically acceptable salt or solvate thereof, can also be used in combination with an anti-osteoporosis agent including a hormonal agent such as raloxifene, or a biphosphonate such as alendronate.

The present invention still further relates to the combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, together with a: (i) tryptase inhibitor; (ii) platelet activating factor (PAF) antagonist; (iii) interleukin converting enzyme (ICE) inhibitor; (iv) IMPDH inhibitor; (v) adhesion molecule inhibitors including VLA-4 antagonist; (vi) cathepsin; (vii) kinase inhibitor such as an inhibitor of tyrosine kinase (such as Btk, Itk, Jak3 or MAP, for example Gefitinib or Imatinib mesylate), a serine / threonine kinase (such as an inhibitor of a MAP kinase such as p38, JNK, protein kinase A, B or C, or IKK), or a kinase involved in cell cycle regulation (such as a cyclin dependent kinase); (viii) glucose-6 phosphate dehydrogenase inhibitor; (ix) kinin-B.sub1. - or B.sub2. -receptor antagonist; (x) anti-gout agent, for example colchicine; (xi) xanthine oxidase inhibitor, for example allopurinol; (xii) uricosuric agent, for example probenecid, sulfinpyrazone or benzbromarone; (xiii) growth hormone secretagogue; (xiv) transforming growth factor (TGF β); (xv) platelet-derived growth factor (PDGF); (xvi) fibroblast growth factor for example basic fibroblast growth factor (bFGF); (xvii) granulocyte macrophage colony stimulating factor (GM-CSF); (xviii) capsaicin cream; (xix) tachykinin NK.sub1. or NK.sub3. receptor antagonist such as NKP-608C, SB-233412 (talnetant) or D-4418; (xx) elastase inhibitor such as UT-77 or ZD-0892; (xxi) TNF-alpha converting enzyme inhibitor (TACE); (xxii) induced nitric oxide synthase (iNOS) inhibitor; (xxiii) chemoattractant receptor-homologous molecule expressed on TH2 cells, (such as a CRTH2 antagonist); (xxiv) inhibitor of P38; (xxv) agent modulating the function of Toll-like receptors (TLR), (xxvi) agent modulating the activity of purinergic receptors such as P2X7; or (xxvii) inhibitor of transcription factor activation such as NFkB, API, or STATS.

A compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, can also be used in combination with an existing therapeutic agent for the treatment of cancer, for example suitable agents include:

- (i) an antiproliferative/antineoplastic drug or a combination thereof, as used in medical oncology, such as an alkylating agent (for example cis-platin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulphan or a nitrosourea); an antimetabolite (for example an antifolate such as a fluoropyrimidine like 5-fluorouracil or tegafur, raltitrexed, methotrexate, cytosine arabinoside, hydroxyurea, gemcitabine or paclitaxel); an antitumour antibiotic (for example an anthracycline such as adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin or mithramycin); an antimitotic agent (for example a vinca alkaloid such as vincristine, vinblastine, vindesine or vinorelbine, or a taxoid such as taxol or taxotere); or a topoisomerase inhibitor (for example an epipodophyllotoxin such as etoposide, teniposide, amsacrine, topotecan or a camptothecin);
- (ii) a cytostatic agent such as an antioestrogen (for example tamoxifen, toremifene, raloxifene, droloxifene or idoxifene), an oestrogen receptor down regulator (for example fulvestrant), an antiandrogen (for example bicalutamide, flutamide, nilutamide or cyproterone acetate), a LHRH antagonist or LHRH agonist (for example goserelin, leuporelin or buserelin), a progestogen (for example megestrol acetate), an aromatase inhibitor (for example as anastrozole, letrozole, vorazole or exemestane) or an inhibitor of 5 α -reductase such as finasteride;
- (iii) an agent which inhibits cancer cell invasion (for example a metalloproteinase inhibitor like marimastat or an inhibitor of urokinase plasminogen activator receptor function);
- (iv) an inhibitor of growth factor function, for example: a growth factor antibody (for example the anti-erbB2 antibody trastuzumab, or the anti-erbB1 antibody cetuximab [C225]), a farnesyl transferase inhibitor, a tyrosine kinase inhibitor or a serine/threonine kinase inhibitor, an inhibitor of the epidermal growth factor family (for example an EGFR family tyrosine kinase inhibitor such as N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, AZD1839), N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) or 6-acrylamido-N-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)quinazolin-4-amine (CI 1033)), an

inhibitor of the platelet-derived growth factor family, or an inhibitor of the hepatocyte growth factor family;

(v) an antiangiogenic agent such as one which inhibits the effects of vascular endothelial growth factor (for example the anti-vascular endothelial cell growth factor antibody bevacizumab, a compound disclosed in WO 97/22596, WO 97/30035, WO 97/32856 or WO 98/13354), or a compound that works by another mechanism (for example linomide, an inhibitor of integrin $\alpha v \beta 3$ function or an angiostatin);

(vi) a vascular damaging agent such as combretastatin A4, or a compound disclosed in WO 99/02166, WO 00/40529, WO 00/41669, WO 01/92224, WO 02/04434 or WO 02/08213;

(vii) an agent used in antisense therapy, for example one directed to one of the targets listed above, such as ISIS 2503, an anti-ras antisense;

(viii) an agent used in a gene therapy approach, for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene-directed enzyme pro-drug therapy) approaches such as those using cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi-drug resistance gene therapy; or

(ix) an agent used in an immunotherapeutic approach, for example ex-vivo and in-vivo approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such as interleukin 2, interleukin 4 or granulocyte-macrophage colony stimulating factor, approaches to decrease T-cell anergy, approaches using transfected immune cells such as cytokine-transfected dendritic cells, approaches using cytokine-transfected tumour cell lines and approaches using anti-idiotypic antibodies.


The invention will now be further explained by reference to the following illustrative examples. In the examples the NMR spectra were measured on a Varian Unity spectrometer at a proton frequency of either 300 or 400 MHz. The MS spectra were measured on either an Agilent 1100 MSD G1946D spectrometer or a Hewlett Packard HP1100 MSD G1946A spectrometer. Preparative HPLC separations were performed using a Waters Symmetry[®] or Xterra[®] column using 0.1% aqueous trifluoroacetic acid:

acetonitrile, 0.1% aqueous ammonia: acetonitrile or 0.1% aqueous ammonium acetate: acetonitrile as the eluant. Microwave reactions were performed in a CEM Discover single

mode microwave. In the following examples all compounds were named using the Chemical Abstracts Service Index Name function within the ACD/Name software package.

5 **Example 1**

4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-4-carboxylic acid




A mixture of 2-chloro-5-iodo-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Prepared
10 as described in WO200144170) (200 mg), 4-carboxyphenylboronic acid (83 mg),
potassium carbonate (140 mg) and dichlorobis(triphenylphosphine)palladium (II) (35 mg)
in 1,4-dioxane (2 mL) / water (1 mL) was heated at 80°C under a nitrogen atmosphere for 3
hours. The products were filtered through diatomaceous earth, washing with methanol (2 x
10 mL). The solvent was removed *in vacuo* and the residue was purified by
15 chromatography (SiO₂, dichloromethane:methanol 97:3 as eluant) and then by
recrystallisation (acetonitrile) to yield the title compound as a colourless solid (150 mg).

MS: APCI(-ve) 422/424 (M-H⁺).

m.p. 267-270°C dec.

20 ¹H NMR (400 MHz, d₆-DMSO) δ 13.03 (1H, s), 8.42 (1H, t), 8.04 (2H, d), 7.84 (2H, d),
7.80 (1H, dd), 7.72 (1H, d), 7.61 (1H, d), 2.98 (2H, d), 1.95 (3H, s), 1.74-1.57 (6H, m),
1.55 (6H, s).

Example 2**4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-3-carboxylic acid****a) [4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid**

Methylolithium (1.6M in diethyl ether, 9.5 mL) was added to a stirred solution of 5-bromo-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]]dec-1-ylmethyl)-benzamide (Prepared as described in WO200144170) (4.8 g) in tetrahydrofuran (120 mL) at -78 °C under an atmosphere of nitrogen. After 10 minutes, triisopropyl borate (15 mL) was added, followed by *tert*-butyllithium (1.7M in pentane, 17 mL). After stirring at -78 °C for 2 hours, saturated aqueous ammonium chloride (125 mL) was cautiously added and the mixture was allowed to warm to room temperature over 16 hours. Ethyl acetate (500 mL) was added and the organic fraction was washed with water (3 x 250 mL) before being dried (MgSO₄), filtered and concentrated *in vacuo* to yield the sub-title compound as a colourless solid (4.2 g).

MS: APCI(+ve) 348/350 (M+H⁺).

b) 4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-3-carboxylic acid

Prepared according to the method of Example 1 using [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a))

(300 mg), 3-iodobenzoic acid (210 mg), potassium carbonate (240 mg) and dichlorobis(triphenylphosphine)palladium (II) (60 mg) to yield the title compound as a solid (40 mg).

5 MS: APCI(-ve) 422/424 ($M-H^+$).


m.p. 255-257°C

1H NMR (400 MHz, d_6 -DMSO) δ 13.15 (1H, s), 8.42 (1H, t), 8.2 (1H, t), 7.99-7.95 (2H, m), 7.77 (1H, dd), 7.68 (1H, d), 7.63 (1H, t), 7.60 (1H, d), 2.97 (2H, d), 1.95 (3H, s), 1.69-1.59 (6H, m), 1.54 (6H, s).

10

Example 3

4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid



15 **a) 4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid, ethyl ester**

Prepared according to the method of Example 1 using [4-chloro-3-

[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (500 mg), ethyl 2-iodobenzoate (600 mg), potassium carbonate (600 mg) and

20 dichlorobis(triphenylphosphine)palladium (II) (50 mg) to give the sub-title compound as a solid (350 mg).

MS: APCI(+ve) 452/454 ($M+H^+$).

b) 4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid


A mixture of 4'-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid, ethyl ester (Example 3 (a)) (350 mg), methanol (2 mL) and aqueous sodium hydroxide (2.5M, 2.0 mL) was heated in a microwave at 70°C for 2 hours. The volatile components were removed *in vacuo* and the residue partitioned between aqueous sodium hydroxide (2M, 20 mL) and dichloromethane (20 mL). The aqueous fraction was acidified to pH 3 with hydrochloric acid (2M) and extracted with ethyl acetate (2 x 20 mL). The combined organics were dried (MgSO₄), filtered and concentrated. Recrystallisation (acetonitrile) gave the title compound as a colourless solid (15 mg). MS: APCI(-ve) 422/424 (M-H⁺).

m.p. 228-230°C

¹H NMR (400 MHz, d₆-DMSO) δ 12.94 (1H, s), 8.38 (1H, t), 7.78 (1H, dd), 7.61 (1H, ddd), 7.51 (1H, d), 7.50 (1H, ddd), 7.41 (1H, dd), 7.38 (1H, dd), 7.34 (1H, d), 2.94 (2H, d), 1.94 (3H, s), 1.70-1.56 (6H, m), 1.53-1.51 (6H, m).

Example 4

2-Chloro-5-[6-(cyanoamino)pyrazinyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



A mixture of [4-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (500 mg), 2,6-dichloropyrazine (900 mg), potassium carbonate (350 mg) and dichlorobis(triphenylphosphine)palladium (II) (90 mg) in 1,4-dioxane (5 mL) / water (2 mL) was heated at 80°C under a nitrogen atmosphere for 1 hour.

The products were filtered through diatomaceous earth, washing with methanol (2 x 10 mL). The solvent was removed *in vacuo* and the residue was purified (SiO₂, dichloromethane:methanol 99:1 as eluant) to give a solid which was taken up in acetonitrile (1.5 mL) and sodium hydrogen cyanamide (270 mg) was added. The mixture
 5 was heated in a microwave at 100°C for 3 hours and then purified (SiO₂, dichloromethane:methanol 95:5 as eluant). Further purification (RP-HPLC, acetonitrile:aqueous trifluoroacetic acid, Symmetry) gave the title compound as a colourless solid (26 mg).


10 MS: APCI(+ve) 422/424 (M+H⁺), 439/441 (M+NH₃+H⁺)

m.p. 215-220°C dec.

¹H NMR (400 MHz, d₆-DMSO) δ 8.41 (1H, t), 8.08 (1H, s), 8.04 (1H, d), 8.00 (1H, s), 7.62 (1H, s), 7.54 (1H, d), 2.97 (2H, d), 1.95 (3H, s), 1.69-1.61 (6H, m), 1.55 (6H, s).

15 Example 5

2-Chloro-5-[3-(cyanamino)pyrazinyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



20 **a) 2-Chloro-5-(3-chloropyrazinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

Prepared according to the method of Example 1, using [4-chloro-3-

[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a))

(1.0 g), 2,3-dichloropyrazine (1.8 g), potassium carbonate (700 mg) and

dichlorobis(triphenylphosphine)palladium (II) (180 mg). Ethyl acetate (100 mL) and water

(50 mL) were added and the organic fraction was washed with hydrochloric acid (2M, 50 mL) and saturated sodium chloride (50 mL) before being dried (MgSO_4) and evaporated. Purification by chromatography (SiO_2 , dichloromethane:methanol 99:1 as eluant) gave the sub-title compound as a solid (750 mg).

MS: APCI(+ve) 416/418 ($\text{M}+\text{H}^+$)

b) 2-Chloro-5-[3-(cyanamino)pyrazinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

A mixture of 2-chloro-5-(3-chloropyrazinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 5 (a)) (340 mg), sodium hydrogen cyanamide (280 mg), tetrabutylammonium bromide (50 mg) and acetonitrile (2 mL) was heated in a microwave at 100°C for 3 hours. Purification (SiO_2 , dichloromethane:methanol: trifluoroacetic acid 96:3:1 as eluant) gave the title compound as a solid (76 mg).


MS: APCI(+ve) 422/424 ($\text{M}+\text{H}^+$)

m.p. 110-115°C dec.

^1H NMR (400 MHz, d_6 -DMSO, 363 K) δ 8.24-8.14 (1H, m), 8.11-8.02 (2H, m), 7.92-7.83 (2H, m), 7.58 (1H, d), 2.98 (2H, d), 1.95 (3H, br s), 1.73-1.58 (6H, m), 1.55 (6H, s).

Example 6

3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-2-pyrazinecarboxylic acid, monosodium salt



a) 2-Chloro-5-(3-cyanopyrazinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

Prepared according to the method of Example 1, using [4-chloro-3-

[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-boronic acid (Example 2 (a))

(0.5 g), 3-chloro-2-pyrazinecarbonitrile (0.6 g), potassium carbonate (350 mg) and
 dichlorobis(triphenylphosphine)palladium (II) (500 mg). The mixture was extracted with
 ethyl acetate (3 x 10 mL) and the combined organic fractions were washed with water (10
 mL) and saturated sodium chloride (10 mL) before being dried (MgSO₄) and evaporated.
 Purification by chromatography (SiO₂, dichloromethane as eluant) gave the sub-title
 compound as a solid (390 mg).

MS: APCI(+ve) 407/409 (M+H⁺)

b) 3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-2-pyrazinecarboxylic acid, monosodium salt

A mixture of 2-chloro-5-(3-cyanopyrazinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 benzamide (Example 6 (a)) (200 mg), methanol (1 mL) and aqueous sodium hydroxide
 (6M, 1.0 mL) was heated in a microwave at 80°C for 2 hours. Purification by
 chromatography (SiO₂, dichloromethane:methanol:trifluoroacetic acid 96:4:1 as eluant),
 redissolution in acetonitrile (2 mL) and formation of the sodium salt by addition of sodium
 hydroxide (1M, 0.47 mL) and filtration of the resulting precipitate gave the title compound
 as a solid (90 mg).


MS: APCI(+ve) 426/428 (M+H⁺).

m.p. 194-195°C dec.

¹H NMR (400 MHz, d₆-DMSO) δ 8.87 (1H, d), 8.71 (1H, d), 8.46 (1H, t), 7.74-7.69 (2H, m), 7.64 (1H, d), 2.95 (2H, d), 1.94 (3H, s), 1.72-1.57 (6H, m), 1.55-1.51 (6H, m).

Example 7

3-[5-Chloro-4-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]-2-pyridinyl]-benzoic acid



Prepared according to the method of Example 1, using 2,5-dichloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-4-pyridinecarboxamide (Prepared as described in WO200144170) (0.5 g), 3-carboxyphenylboronic acid (0.5 g), potassium carbonate (350 mg) and dichlorobis(triphenylphosphine)palladium (II) (50 mg). Purification (SiO₂, dichloromethane:methanol 97:3 as eluant) and then by recrystallisation (acetonitrile:diethyl ether) gave the title compound as a solid (24 mg).


MS: APCI(+ve) 425/427 (M+H⁺).

m.p. 200-202°C

¹H NMR (400 MHz, d₆-DMSO) δ 8.80 (1H, s), 8.69 (1H, s), 8.60 (1H, t), 8.36 (1H, d), 8.06 (1H, s), 8.03 (1H, d), 7.65 (1H, dd), 2.99 (2H, d), 1.96 (3H, s), 1.72-1.58 (6H, m), 1.58-1.53 (6H, m).

Example 8

2-Chloro-5-[3-[(methylsulfonyl)amino]pyrazinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



A mixture of 2-chloro-5-(3-chloropyrazinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 5 (a)) (340 mg), methanesulfonamide (70 mg), potassium *tert*-butoxide (28 mg) and acetonitrile (1 mL) was heated in a microwave at 140°C for 4 hours.

5 Purification (Varian NH₂ cartridge using methanol (100 mL) and then 1 % trifluoroacetic acid in methanol (100 mL) as eluant, then RP-HPLC, acetonitrile:aqueous ammonium acetate, Symmetry) gave the title compound as a solid (9 mg).


MS: APCI(+ve) 475/477 (M+H⁺)

10 m.p. 245°C dec.

¹H NMR (400 MHz, d₆-DMSO) δ 8.52-8.39 (2H, m), 8.36 (1H, t), 7.84-7.76 (2H, m), 7.66-7.60 (1H, m), 3.34 (s, 3H), 2.96 (2H, d), 1.93 (3H, s), 1.70-1.57 (6H, m), 1.52 (6H, s).

Example 9

15 **2-Chloro-5-[3-(1-*H*-tetrazol-5-yl)pyrazinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**



A mixture of 2-chloro-5-(3-cyanopyrazinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 6 (a)) (180 mg), azidotrimethylsilane (0.15 mL) and dibutyltin oxide (20 mg) in toluene (1 mL) was heated at 95°C for 16 hours. Further azidotrimethylsilane (0.15 mL) was added and the mixture was heated at 95°C for 8 hours. Purification (Varian NH₂ cartridge using methanol (100 mL) and then 1 % trifluoroacetic acid in methanol (100 mL) as eluant), subsequent recrystallisation (methanol) and washing of the resulting solid with *isohexane* gave the title compound as a solid (55 mg).


MS: APCI(+ve) 450/452 (M+H⁺)

m.p. 254°C dec.

¹H NMR (400 MHz, d₆-DMSO) δ 8.96 (1H, d), 8.90 (1H, d), 8.41 (1H, t), 7.57-7.46 (3H, m), 2.93 (2H, d), 1.95 (3H, s), 1.71-1.56 (6H, m), 1.54-1.47 (6H, m).

Example 10

2-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-3-pyridinecarboxylic acid



a) 2-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-3-pyridinecarboxylic acid, methyl ester

A mixture of [4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-
 5 boronic acid (Prepared as described in Example 2 (a)) (200 mg), 2-chloronicotinic acid,
 methyl ester (118 mg), potassium carbonate (166 mg) and
 tetrakis(triphenylphosphine)palladium(0) (69 mg) in tetrahydrofuran (2 mL) / water (1 mL)
 was heated at 65°C under a nitrogen atmosphere for 1 hour. The products were filtered
 through diatomaceous earth, washing with methanol (2 x 10 mL). The solvent was
 10 removed *in vacuo* and the residue was purified (SiO₂, dichloromethane:methanol 99:1 as
 eluant) to yield the sub-title compound as a solid (200 mg).
 MS: APCI(+ve) 439/441 (M+H⁺).

b) 2-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-3-pyridinecarboxylic acid

Potassium hydroxide (100 mg) in methanol (2 mL) was added to a stirred solution of 2-[4-
 chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-3-
 pyridinecarboxylic acid, methyl ester (Example 10 (a)) (200 mg) in methanol (1 mL). The
 20 reaction was stirred at room temperature for 24 hours then evaporated. The residue was
 dissolved in water (5 mL) and the solution was acidified to pH 5 with 2 M aqueous
 hydrochloric acid. The resulting solid was collected by filtration and purified (Varian NH₂

cartridge using methanol (100 mL) and then 5 % acetic acid in methanol (100 mL) as eluent) to afford the title compound as a solid (25 mg).


MS: APCI(-ve) 423/425 ($M-H^+$).

5 m.p. 169-172°C.

1H NMR (400 MHz, d_6 -DMSO) δ 8.76 (1H, dd), 8.43 (1H, t), 8.13 (1H, dd), 7.62-7.47 (4H, m), 2.95 (2H, d), 1.94 (3H, s), 1.67 (3H, d), 1.59 (3H, d), 1.52 (6H, s).

10 **Example 11**

5-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid



15 **a) 5-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid, methyl ester**

Prepared according to the method of Example 10 (a) using [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Prepared as described in Example 2 (a)) (200 mg) and 5-bromo-3-pyridinecarboxylic acid methyl ester (150 mg). Purification (SiO_2 , dichloromethane:methanol 99:1 as eluant) afforded the sub-

20 title compound as a solid (100 mg).

MS: APCI(+ve) 439/441 ($M+H^+$).

b) 5-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid

Prepared according to the method of Example 10 (b) using 5-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid, methyl ester (Example 11 (a)) (100 mg). Purification (Varian NH₂ cartridge using methanol:dichloromethane 1:1(100 mL) and then acetic acid:methanol:dichloromethane 1:10:10 (100 mL) as eluant) afforded the title compound as a solid (33 mg).


MS: APCI(+ve) 425/427 (M+H⁺).

m.p. 178-182°C.

¹H NMR (400 MHz, d₆-DMSO) δ 9.14 (1H, d), 9.08 (1H, d), 8.50 (1H, t), 8.42 (1H, t), 7.88 (1H, d), 7.85 (1H, d), 7.81 (1H, d), 2.98 (2H, d), 1.95 (3H, s), 1.68 (3H, d), 1.61 (3H, d), 1.55 (6H, s).

Example 12

2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-pyridinecarboxylic acid



a) 2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-pyridinecarboxylic acid, 1,1-dimethylethyl ester

Prepared according to the method of Example 10 (a) using [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Prepared as described in Example 2 (a)) (200 mg) and 2-chloro-4-pyridinecarboxylic acid, 1,1-

dimethylethyl ester (150 mg). Purification (SiO₂, 99:1 dichloromethane:methanol 99:1 as eluant) afforded the sub-title compound as a solid (150 mg).

MS: APCI(+ve) 481/483 (M+H⁺).

b) 2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-pyridinecarboxylic acid

Trifluoroacetic acid (1 mL) was added to a stirred solution of 2-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-pyridinecarboxylic acid 1,1-dimethylethyl ester (Example 12 (a)) (150 mg) in dichloromethane (3 mL). The reaction was stirred at room temperature overnight then evaporated. Purification (SiO₂, dichloromethane:methanol 99:1 as eluant) afforded the title compound as a solid (17 mg).


MS: APCI(-ve) 423/425 (M-H⁺).

m.p. 284-287°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.82 (1H, d), 8.45 (1H, t), 8.34 (1H, s), 8.16 (1H, dd), 8.11 (1H, d), 7.80 (1H, d), 7.62 (1H, d), 2.98 (2H, d), 1.95 (3H, s), 1.68 (3H, d), 1.61 (3H, d), 1.55 (6H, s).

Example 13

2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-6-methyl-3-pyridinecarboxylic acid



a) 2-Chloro-6-methyl-3-pyridinecarboxylic acid, methyl ester

To a stirred solution of 2-chloro-6-methyl-3-pyridinecarboxylic acid (200 mg) in dichloromethane (2 mL) at 0°C under nitrogen was added *N,N*-dimethylformamide (1 drop) and oxalyl chloride (0.3 mL). The reaction mixture was stirred at room temperature for 30 minutes, then evaporated to dryness and redissolved in dichloromethane (2 mL).

5 Methanol (2mL) was added dropwise and the mixture was stirred at room temperature for 10 minutes before being evaporated to give the sub-title compound as a solid (210 mg).

MS: APCI(+ve) 186/188 ($M+H^+$).

1H NMR (400 MHz, d_6 -DMSO) δ 8.16 (1H, d), 7.42 (1H, d), 3.87 (3H, s), 2.52 (3H, s).

10 **b) 2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-6-methyl-3-pyridinecarboxylic acid, methyl ester**

Prepared according to the method of Example 10 (a) using [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Prepared as described in Example 2 (a)) (200 mg) and 2-chloro-6-methyl-3-pyridinecarboxylic acid methyl ester (Example 13 (a)) (133 mg). Purification (SiO_2 , dichloromethane: methanol 99:1 as eluant) afforded the sub-title compound as a solid (130 mg).

MS: APCI(+ve) 453/455 ($M+H^+$).

20 **c) 2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-6-methyl-3-pyridinecarboxylic acid**

Prepared according to the method of Example 10 (b) using 2-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-6-methyl-3-pyridinecarboxylic acid, methyl ester (Example 13 (b)) (130 mg). Purification (Varian NH_2 cartridge using methanol:dichloromethane 1:1(100 mL) and then acetic acid:methanol:dichloromethane 1:10:10 (100 mL) as eluant) afforded the title compound as a solid (27 mg).


30 MS: APCI(-ve) 437/439 ($M-H^+$).

m.p. 209-211°C.

¹H NMR (400 MHz, CD₃OD) δ 7.98 (1H, d), 7.66 - 7.61 (2H, m), 7.51 (1H, d), 7.31 (1H, d), 3.07 (2H, s), 2.59 (3H, s), 1.98 (3H, s), 1.77 (3H, d), 1.70 (3H, d), 1.63 (6H, s).

Example 14

(2*S*)-2-[[4'-chloro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid



a) 4-Chloro-2'-hydroxy-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-[1,1'-biphenyl]-3-carboxamide

A mixture of 2-chloro-5-iodo-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Prepared as described in WO200144170) (1 g), (2-hydroxyphenyl)-boronic acid (450 mg), potassium carbonate (700 mg) and dichlorobis(triphenylphosphine)palladium(II) (175 mg) in tetrahydrofuran (10 mL) / water (10 mL) was heated at 50°C under a nitrogen atmosphere for 1 hour. The mixture was concentrated and then extracted with dichloromethane (3 x 50 mL). The combined organic extracts were filtered through diatomaceous earth and evaporated. Purification (SiO₂, dichloromethane:methanol 99:1 as eluant) afforded the sub-title compound as a solid (900 mg).

MS: APCI(+ve) 396/398 (M+H⁺).

b) (2*S*)-2-[[4'-chloro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid, methyl ester

Methyl (*R*)- 2-chloropropionate (311 mg) was added to a stirred solution of 4-chloro-2'-hydroxy-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-[1,1'-biphenyl]-3-carboxamide (Example 14 (a)) (250 mg) and potassium carbonate (174 mg) in acetone (4 mL). The reaction mixture was heated to 55 °C under a nitrogen atmosphere for 10 hours, was then allowed to cool and was concentrated. The residue was partitioned between water (20 mL) and dichloromethane (20 mL). The layers were separated and the aqueous was extracted with dichloromethane (2 x 20 mL). The combined organics were dried, filtered and evaporated. Purification (SiO₂, dichloromethane as eluant) afforded the sub-title compound as a solid (110 mg).

MS: APCI(+ve) 482/484 (M+H⁺).

¹H NMR (400 MHz, d₆-DMSO) δ 8.39 (1H, t), 7.67-7.60 (2H, m), 7.52 (1H, d), 7.39-7.29 (2H, m), 7.07 (1H, t), 6.96 (1H, d), 5.10 (1H, q), 3.67 (3H, s), 3.00-2.89 (2H, m), 1.93 (3H, s), 1.67 (3H, d), 1.59 (3H, d), 1.53 (6H, s), 1.45 (3H, d).

c) (2*S*)-2-[[4'-chloro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid

Prepared according to the method of Example 10 (b) using (2*S*)-2-[[4'-chloro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid, methyl ester (Example 14 (b)) (110 mg). The residue was dissolved in deionised water (5 mL) and the solution was acidified to pH 5 with 2 M aqueous hydrochloric acid. The resulting solid was collected by filtration and washed with water to give the title compound as a solid (75 mg).


MS: APCI(+ve) 468/470 (M+H⁺).

m.p. 107-110°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.39 (1H, t), 7.70-7.63 (2H, m), 7.51 (1H, d), 7.38-7.29 (2H, m), 7.05 (1H, t), 6.93 (1H, d), 4.93 (1H, q), 2.94 (2H, d), 1.93 (3H, s), 1.66 (3H, d), 1.59 (3H, d), 1.52 (6H, s), 1.45 (3H, d).

Example 15

[[4'-Chloro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-acetic acid



a) [[4'-Chloro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-acetic acid, ethyl ester

Ethyl chloroacetate (300 mg) was added to a stirred solution of 4-chloro-2'-hydroxy-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-[1,1'-biphenyl]-3-carboxamide (Example 14 (a)) (250 mg) and potassium carbonate (174 mg) in acetone (4 mL). The reaction mixture was heated to 55 °C under a nitrogen atmosphere for 1 hour, then allowed to cool and concentrated. The residue was partitioned between deionised water (20 mL) and dichloromethane (20 mL). The layers were separated and the aqueous fraction was extracted with dichloromethane (2 x 20 mL). The combined organics were dried, filtered and evaporated. Recrystallisation from acetonitrile gave the sub-title compound (140 mg).

MS: APCI(+ve) 482/484 (M+H⁺).

¹H NMR (400 MHz, d₆-DMSO) δ 8.35 (1H, t), 7.63 (1H, dd), 7.58 (1H, d), 7.51 (1H, d), 7.39-7.30 (2H, m), 7.12-7.01 (2H, m), 4.83 (2H, s), 4.15 (2H, q), 2.94 (2H, d), 1.93 (3H, s), 1.67 (3H, d), 1.59 (3H, d), 1.52 (6H, s), 1.20 (3H, t).

b) [[4'-Chloro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-acetic acid

A solution of potassium hydroxide (100 mg) in water (2 mL) was added to a solution of [[4'-chloro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-acetic acid, ethyl ester (Example 15 (a)) (140 mg) in methanol (2 mL) in a 10 mL

microwave vial and this was heated at 70°C for 5 minutes within a microwave. The reaction mixture was concentrated and then acidified to pH 5 using 2M aqueous hydrochloric acid. The solid was collected by filtration and purification (Varian NH₂ cartridge using methanol:dichloromethane 1:1(100 mL) and then acetic acid:methanol:dichloromethane 1:10:10 (100 mL) as eluant) gave the title compound as a solid (55 mg).


MS: APCI(+ve) 454/456 (M+H⁺).

m.p. 208-210°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.36 (1H, t), 7.64 (1H, dd), 7.57 (1H, d), 7.51 (1H, d), 7.38-7.30 (2H, m), 7.06 (1H, t), 7.00 (1H, d), 4.70 (2H, s), 2.94 (2H, d), 1.93 (3H, s), 1.66 (3H, d), 1.59 (3H, d), 1.53 (6H, s).

Example 16

3-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid



To a solution of 4-chloro-2'-hydroxy-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-[1,1'-biphenyl]-3-carboxamide (Prepared as described in Example 14 (a)) (200 mg) and potassium *tert*-butoxide (120 mg) in tetrahydrofuran (2 mL) was added β-propiolactone (0.06 mL). The reaction was stirred at room temperature under a nitrogen atmosphere for 16 hours then evaporated. The residue was suspended in deionised water (5 mL) and acidified to pH 5 with 2 M aqueous hydrochloric acid. The solid was collected by filtration and purified (Varian NH₂ cartridge using methanol (100 mL) and then 5 % acetic acid in

methanol (100 mL) as eluant). Further purification (SiO₂, dichloromethane:methanol 99:1 as eluant) afforded the title compound as a solid (10 mg).


MS: APCI(+ve) 468/470 (M+H⁺).

5 m.p. 163-164°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.36 (1H, t), 7.57 (1H, dd), 7.47-7.42 (2H, m), 7.40-7.33 (2H, m), 7.15 (1H, d), 7.06 (1H, t), 4.21 (2H, t), 2.95 (2H, d), 2.64 (2H, t), 1.94 (3H, s), 1.67 (3H, d), 1.59 (3H, d), 1.53 (6H, s).

10 **Example 17**

5-Chloro-2-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino] carbonyl]phenyl]-3-pyridinecarboxylic acid



a) 2,5-Dichloro-3-pyridinecarboxylic acid, methyl ester

15 2,5-Dichloro-3-pyridinecarbonyl chloride (1.0 g) was added dropwise over 10 minutes to methanol (10 ml) with stirring under a nitrogen atmosphere. The resulting solution was stirred at room temperature for 16 hours and concentrated *in vacuo* to yield the sub-title compound as a colourless solid (0.95 g)

20 MS: APCI(+ve) 206 (M+H⁺).

b) 5-Chloro-2-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid, methyl ester

Prepared according to the method of Example 1 using [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (300 mg), 2,5-dichloro-3-pyridinecarboxylic acid, methyl ester (Example 17 (a)) (180 mg), potassium carbonate (240 mg) and dichlorobis(triphenylphosphine)palladium (II) (50 mg) to give the sub-title compound as a colourless solid (350 mg).

MS: APCI(+ve) 475 (M+H⁺).

c) 5-Chloro-2-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid

Prepared according to the method of Example 10 (b) using 5-chloro-2-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid, methyl ester (Example 17 (b)) (350 mg). Purification (SiO₂, dichloromethane:methanol 99:1, then dichloromethane:methanol:trifluoroacetic acid 98:1:1 as eluant) gave the title compound as a colourless solid (32 mg).


MS: APCI(+ve) 459 (M+H⁺).

m.p. 165-168°C.

¹H NMR (300 MHz, d₆-DMSO) δ 8.85 (1H, t), 8.44 (1H, t), 8.28 (1H, d), 7.61-7.55 (3H, m), 2.95 (2H, d), 1.94 (3H, s), 1.72-1.55 (6H, m), 1.52 (6H, s).

Example 18

4'-Chloro-6-methyl-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid



a) 2-Iodo-3-methyl-benzoic acid, methyl ester

Prepared according to the method of Example 13 (a) using 2-iodo-3-methyl-benzoic acid (1.0 g) to yield the sub-title compound as an oil (0.94g).

5

MS: APCI(+ve) 276 ($M+H^+$).

b) 4'-Chloro-6-methyl-3'--[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]- [1,1'-biphenyl]-2-carboxylic acid, methyl ester

10

Prepared according to the method of Example 1 using [4-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (300 mg), 2-iodo-3-methyl-benzoic acid, methyl ester (Example 18 (a)) (250 mg), potassium carbonate (240 mg) and dichlorobis(triphenylphosphine)palladium (II) (50 mg) to give the sub-title compound as a solid (250 mg).

15

MS: APCI(+ve) 452 ($M+H^+$).

c) 4'-Chloro-6-methyl-3'--[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]- [1,1'-biphenyl]-2-carboxylic acid

20

Prepared according to the method of Example 10 (b) using 4'-Chloro-6-methyl-3'--[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]- [1,1'-biphenyl]-2-carboxylic acid, methyl ester (Example 18 (b)) (250 mg). Purification (SiO_2 , dichloromethane, then dichloromethane:methanol:trifluoroacetic acid 98:2:1 as eluant) gave the title compound as a colourless solid (42 mg).


MS: APCI(+ve) 438 (M+H⁺).

m.p. 151-154°C.

¹H NMR (400 MHz, d₆-DMSO) δ 12.60 (1H, s), 8.31 (1H, t), 7.61 (1H, d), 7.49 (1H, d),
 5 7.48 (1H, d), 7.38 (1H, dd), 7.22 (1H, dd), 7.13 (1H, d), 2.93 (2H, d), 2.08 (3H, s), 1.93
 (3H, s), 1.70-1.55 (6H, m), 1.52 (6H, s).

Example 19

3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-
 10 thiophenecarboxylic acid



a) 3-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-
 thiophenecarboxylic acid, methyl ester

Prepared according to the method of Example 1, using [4-chloro-3-

15 [[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a))
 (0.3 g), 3-bromo-2-thiophenecarboxylic acid, methyl ester (0.19 g), potassium carbonate
 (240 mg) and dichlorobis(triphenylphosphine)palladium (II) (25 mg). The mixture was
 filtered through diatomaceous earth, washing with methanol (2 x 30 ml) and was
 concentrated *in vacuo*. The residue was partitioned between dichloromethane (30 ml) and
 20 deionised water (10 ml), the layers were separated and the aqueous fraction was extracted
 with dichloromethane (2 x 30 ml). The combined organic layers were dried (MgSO₄),
 evaporated and purified (SiO₂, 85:15 isohexane:ethyl acetate to give the sub-title
 compound as a colourless solid (160 mg).

MS: APCI(+ve) 444 ($M+H^+$).

b) 3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-thiophenecarboxylic acid

- 5 Prepared according to the method of Example 10 (b) using 3-[4-chloro-3-
[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-thiophenecarboxylic acid,
methyl ester (Example 19 (a)) (160 mg). Purification (SiO_2 , dichloromethane, then
dichloromethane:methanol:trifluoroacetic acid 98:2:1 as eluant, then RP-HPLC,
acetonitrile:aqueous ammonium acetate, Symmetry) gave the title compound as a
10 colourless solid (47 mg).


MS: APCI(+ve) 430 ($M+H^+$).

m.p. 140-143°C.

- ¹H NMR (400 MHz, d_6 -DMSO) δ 8.34 (1H, t), 7.71 (1H, d), 7.59-7.54 (1H, m), 7.49-7.43
15 (2H, m), 7.15 (1H, d), 2.94 (2H, d), 1.93 (3H, s), 1.70-1.56 (6H, m), 1.52 (6H, s).

Example 20

6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid



- 20 a) 6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid, methyl ester

Prepared according to the method of Example 10 (a) using [4-chloro-3-
[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a))
(250 mg) and 6-chloro-2-pyridinecarboxylic acid, methyl ester (150 mg). Purification
(SiO₂, ethyl acetate:*isohexane* 25:75 as eluent) afforded the sub-title compound as a solid
5 (120 mg).

¹H NMR (400 MHz, d₆-DMSO) δ 8.46 (1H, t), 8.32 (1H, dd), 8.22 - 8.16 (2H, m), 8.12
(1H, t), 8.05 (1H, dd), 7.65 (1H, d), 3.92 (3H, s), 2.98 (2H, d), 1.96 (3H, s), 1.69 (3H, d),
1.62 (3H, d), 1.56 (6H, s).

10 MS: APCI(+ve) 439/441 (M+H⁺).

b) 6-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid

Prepared according to the method of Example 10 (b) using 6-[4-chloro-3-

15. [[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid
methyl ester (Example 20 (a)) (120 mg). Purification (Varian NH₂ cartridge using methanol
(100 mL) and then acetic acid:methanol 1:20 (100 mL) as eluant) afforded the title
compound as a solid (65 mg).


20 ¹H NMR (400 MHz, d₆-DMSO) δ 8.45 (1H, s), 8.26 - 8.14 (3H, m), 8.03 (1H, t), 7.96 (1H,
d), 7.62 (1H, d), 2.99 (2H, d), 1.95 (3H, s), 1.75 - 1.48 (12H, m).

MS: APCI(+ve) 425/427 (M+H⁺).

m.p. 170-173°C.

25 **Example 21**

3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid



a) **2-Chloro-5-(2-chloro-3-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**
 [4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid
 (Example 2 (a)) (3.1 g) was added to a solution of 3-bromo-2-chloropyridine (1.7 g) in
 5 tetrahydrofuran (30 mL). A solution of potassium carbonate (2.4 g) in water (30 mL) was
 added followed by dichlorobis(triphenylphosphine)palladium (II) (0.25 g) and the mixture
 was stirred under a nitrogen atmosphere at room temperature for 3 hours. The reaction
 mixture was concentrated and the residue partitioned between dichloromethane and water.
 The layers were separated and the organic fraction was dried (MgSO₄), filtered and
 10 concentrated. Purification by chromatography (SiO₂, 70:30 *isohexane*:ethyl acetate, then
 60:40 *isohexane*:ethyl acetate as eluant) gave the sub-title compound as a colourless solid
 (1.85g)

MS: APCI(+ve) 415/417 (M+H⁺).

b) **2-Chloro-5-(2-cyano-3-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**
 A mixture of 2-chloro-5-(2-chloro-3-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 benzamide (Example 21 (a)) (0.25 g), bis(dibenzylideneacetone)palladium (0.12 g), 1,1'-
 bis(diphenylphosphino)ferrocene (0.29 g) and copper (I) cyanide (0.34 g) in 1,4-dioxane (3
 20 mL) was heated at 130°C in a microwave for 2 hours. The reaction was concentrated to
 dryness and partitioned between dichloromethane and water. The organic fraction was
 dried (MgSO₄), filtered and concentrated to dryness before being purified by
 chromatography (SiO₂, 1:1 ethyl acetate: *isohexane* as eluant) to give the sub-title
 compound as a foam (0.20 g)

MS: APCI(+ve) 406/408 ($M+H^+$).

c) 3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid


A mixture of 2-chloro-5-(2-cyano-3-pyridinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.20 g) (Example 21 (b)) and potassium hydroxide (0.17 g) in water (1 mL) and ethanol (10 mL) was heated at reflux for 16 hours. The reaction mixture was then concentrated to dryness and the residue redissolved in water (2 mL) and concentrated hydrochloric acid (2 mL). This mixture was then heated at reflux for 5 hours, allowed to cool to room temperature and was purified by RP-HPLC (acetonitrile:aqueous ammonium acetate, Symmetry) to give the title compound as a solid (99 mg).

MS: APCI(+ve) 425/427 ($M+H^+$).

¹H NMR (400 MHz, d₆-DMSO) δ 8.38 - 8.33 (2H, m), 7.72 - 7.65 (2H, m), 7.57 (1H, d), 7.46 (1H, d), 7.24 (1H, dd), 2.95 (2H, d), 1.94 (3H, s), 1.66 (3H, d), 1.60 (3H, d), 1.53 (6H, s).

Example 22

2-Chloro-5-[2-(1*H*-tetrazol-5-yl)-3-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



Prepared according to the method of Example 9 using 2-chloro-5-(2-cyano-3-pyridinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 21 (b)) (0.13 g). Purification


(Varian NH₂ cartridge using methanol and then 1 % trifluoroacetic acid in methanol as eluant) followed by RP-HPLC (acetonitrile:aqueous trifluoroacetic acid, Symmetry) gave the title compound as a solid (36 mg).

5 MS: APCI(+ve) 449 (M+H⁺).

¹H NMR (d₆-DMSO, 300MHz) δ 8.84-8.82 (1H, d), 8.36-8.32 (1H, t), 8.02-7.99 (1H, d), 7.76-7.72 (1H, m), 7.49-7.46 (1H, d), 7.36-7.29 (2H, m), 2.93-2.91 (2H, d), 1.93 (3H, s), 1.68-1.57 (6H, q), 1.50 (6H, s).

10 **Example 23**

2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-oxazolecarboxylic acid



a) 4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-benzoic acid

15 Potassium peroxomonosulfate (1.9 g) was added to a solution of 2-chloro-5-formyl-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Prepared as described in WO200144170) (1.0 g) in *N,N*-dimethylformamide (15 mL) and the mixture was stirred at room temperature for 5 hours. Ethyl acetate (50 mL) and aqueous hydrochloric acid (20 mL, 1M) were added, the layers were separated and the aqueous fraction was extracted with ethyl acetate (50 mL). The combined organics were washed with water (30 mL) and saturated aqueous sodium chloride (30 mL) before being dried (MgSO₄), filtered and concentrated to give the sub-title compound as a colourless solid (1.0 g).

20

MS: APCI(+ve) 348/350 ($M+H^+$).

b) *N*-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]benzoyl]-L-serine, methyl ester

1-Hydroxybenzotriazole (0.88 g) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.1 g) were added to a stirred solution of 4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-benzoic acid (Example 23 (a)) (1.0 g) and triethylamine (1.6 mL) in dichloromethane (15 mL) under nitrogen at 0°C. L-Serine methyl ester hydrochloride (0.90 g) was added and the mixture was allowed to warm to room temperature over 16 hours. Water (20 mL) was added, the layers were separated and the aqueous fraction was extracted with dichloromethane (2 x 20 mL). The combined organic layers were washed with aqueous hydrochloric acid (20 mL, 2M) and saturated aqueous sodium chloride (20 mL) before being dried ($MgSO_4$), filtered and concentrated to yield the sub-title compound as a colourless solid (1.1 g).

MS: APCI(+ve) 449/451 ($M+H^+$).

c) 2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-oxazolecarboxylic acid, methyl ester

Diethylaminosulfur trifluoride (0.41 mL) was added dropwise over 10 minutes to a stirred solution of *N*-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]benzoyl]-L-serine, methyl ester (Example 23 (b)) (1.0 g) in dichloromethane (15 mL) at -78°C under nitrogen. The mixture was stirred at -78°C for 1 hour and was then allowed to warm to -30°C. Saturated aqueous sodium hydrogen carbonate (5 mL) was added, the mixture was allowed to warm to room temperature and the layers were separated. The aqueous fraction was extracted with dichloromethane (2 x 10 mL), the combined organic layers were combined and then dried ($MgSO_4$), filtered and concentrated before being purified by chromatography (SiO_2 , dichloromethane as eluant). The material was dissolved in dichloromethane (15 mL), cooled to 0°C and bromotrichloromethane (0.44 mL) was added dropwise. The mixture was stirred under nitrogen for 15 minutes, 1,8-diazabicyclo[5.4.0]undec-7-ene (0.67 mL) was added dropwise and the mixture was allowed to warm to room temperature over 16 hours. Water (10 mL) was added, the layers

separated and the aqueous fraction was extracted with dichloromethane (2 x 15 mL). The combined organic layers were washed with aqueous hydrochloric acid (10 mL, 2M) and saturated aqueous sodium chloride (10 mL) before being dried (MgSO_4), filtered and concentrated to leave the sub-title product (0.91 g).

5

MS: APCI(+ve) 429 ($\text{M}+\text{H}^+$).

d) 2-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-oxazolecarboxylic acid

10 Sodium hydroxide (75 mg) in water (2 mL) was added to a stirred solution of 2-[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-oxazolecarboxylic acid, methyl ester (Example 23 (c)) (400 mg) in tetrahydrofuran (5 mL). The reaction was stirred at room temperature for 24 hours. The mixture was concentrated, acidified with aqueous hydrochloric acid (2M) and the resulting precipitate was collected by filtration,
15 washing with water (2 x 10 mL) and methanol (10 mL) to give the title compound as a solid (86 mg).


MS: APCI(+ve) 415/417 ($\text{M}+\text{H}^+$).

m.p. 254-257°C.

20 ^1H NMR (400 MHz, d_6 -DMSO) δ 8.62 (1H, s), 8.48 (1H, s), 8.02 (1H, dd), 7.99 (1H, d), 7.67 (1H, d) 2.99 (2H, d), 1.91 (3H, m), 1.73-1.57 (6H, m), 1.57-1.51 (6H, m).

Example 24

25 **4'-Chloro-4-methyl-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid**



a) 2-Iodo-5-methyl-benzoic acid, methyl ester

Prepared according to the method of Example 13 (a) using 2-iodo-5-methyl-benzoic acid
 5 (1.0 g) to give the sub-title compound as a colourless oil (1.0 g)

MS: APCI(+ve) 276 ($M+H^+$).

**b) 4'-Chloro-5-methyl-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-
 10 biphenyl]-3-carboxylic acid, methyl ester**

Prepared according to the method of Example 14 (a) using [4-chloro-3-
 [[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a))
 (300 mg) and 2-iodo-5-methyl-benzoic acid, methyl ester (Example 24 (a)) (240 mg) at
 room temperature. Purification (SiO_2 , ethyl acetate:isohexane 15:85 as eluent) afforded the
 15 sub-title compound as a colourless solid (160 mg).

MS: APCI(+ve) 452 ($M+H^+$).

**c) 4'-Chloro-4-methyl-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-
 20 biphenyl]-2-carboxylic acid**

Prepared according to the method of Example 23 (d), using 4'-chloro-5-methyl-3'-
 [[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-3-carboxylic acid,
 methyl ester (Example 24 (b)) (160 mg). The mixture was concentrated and acidified with

aqueous hydrochloric acid (2M) and the residue was purified firstly by chromatography (SiO₂, dichloromethane, then 99:1 dichloromethane:methanol as eluant), and then by recrystallisation from acetonitrile:water to give the title compound as a solid (24 mg).


MS: APCI(+ve) 438/440 (M+H⁺).

m.p. 214-216°C.

¹H NMR (400 MHz, d₆-DMSO) δ 12.88 (1H, s), 8.37 (1H, t), 7.59 (1H, d), 7.49 (1H, d), 7.42 (1H, dd), 7.34 (1H, dd), 7.31 (1H, d), 7.30 (1H, d), 2.94 (2H, d), 2.38 (3H, s), 1.94 (3H, s), 1.70-1.56 (6H, m), 1.52 (6H, d).

Example 25

6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-N-(methylsulfonyl)-2-pyridinecarboxamide



To a stirred mixture of 6-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid (Example 20) (250 mg) and methanesulfonamide (57 mg) in dichloromethane (3 mL) was added 4-(dimethylamino)pyridine (73 mg) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (115 mg). The reaction was stirred at room temperature under nitrogen for 16 hours. Dichloromethane (100 mL) was then added and the solution was washed with water (50 mL) and aqueous hydrochloric acid (2 x 50 mL, 2M) before being dried (MgSO₄), filtered and evaporated. Purification by chromatography (SiO₂, dichloromethane:methanol 98:2 as eluant) and then further purification (Varian NH₂

cartridge using 1:1 dichloromethane:methanol (100 mL) and then 1:10:10 acetic acid:dichloromethane:methanol (100 mL) as eluant) afforded the title compound as a solid (190 mg).

5 MS: APCI(+ve) 502/504 ($M+H^+$).


m.p. 160-162°C.

1H NMR (400 MHz, d_6 -DMSO) δ 8.46 (1H, t), 8.42-8.36 (2H, m), 8.32 (1H, d), 8.14 (1H, t), 8.05 (1H, d), 7.64 (1H, d), 3.38 (3H, s), 2.99 (2H, d), 1.96 (3H, s), 1.68 (3H, d), 1.61 (3H, d), 1.56 (6H, s).

10

Example 26

N-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-glycine



15 a) 5-(2-Amino-3-pyridinyl)-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

To a stirred mixture of [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (1.0 g) and 2-amino-3-bromopyridine (640 mg) in toluene (10 mL) and ethanol (10 mL) was added a solution of sodium carbonate (610 mg) in water (10 mL), followed by

20 tetrakis(triphenylphosphine)palladium(0) (140 mg). The mixture was heated to 50°C for 2 hours, then allowed to cool and was concentrated. The residue was partitioned between dichloromethane (100 mL) and water (100 mL), the layers were separated and the aqueous was extracted with dichloromethane (2 x 50 mL). The combined organics were evaporated.

Purification by chromatography (SiO₂, 3:1 *iso*hexane:ethyl acetate as the eluant) gave the sub-title compound as a solid (500 mg).

MS: APCI(+ve) 396/398 (M+H⁺).

¹H NMR (400 MHz, d₆-DMSO) δ 8.31 (1H, t), 7.98 (1H, dd), 7.66-7.52 (1H, m), 7.48 - 7.44 (2H, m), 7.37 (1H, dd), 6.67 (1H, dd), 5.77 (2H, s), 2.96 (2H, d), 1.94 (3H, s), 1.67 (3H, d), 1.60 (3H, d), 1.52 (6H, s).

b) *N*-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-glycine 1,1-dimethylethyl ester

5-(2-Amino-3-pyridinyl)-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 26 (a)) (300 mg), *tert*-butylbromoacetate (0.5 mL) and acetonitrile (2 mL) were placed in a 10 mL vial and heated at 70°C for 2 hours in a microwave. The mixture was then concentrated and subsequent purification by chromatography (SiO₂, dichloromethane:methanol 95:5 as eluant) afforded the sub-title compound as a solid (200 mg).

MS: APCI(+ve) 510/512 (M+H⁺).

c) *N*-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-glycine

To a solution of *N*-[3-[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-glycine 1,1-dimethylethyl ester (Example 26 (b)) (200 mg) in dichloromethane (3 mL) was added trifluoroacetic acid (1 mL). The mixture was stirred at room temperature for 2 hours and then poured into saturated aqueous sodium bicarbonate (100 mL). This was washed with dichloromethane (3 x 50 mL) and the aqueous fraction was filtered to give a solid which was purified by chromatography (SiO₂, dichloromethane: methanol 90:10 as eluant) to afford the title compound as a solid (40 mg).

MS: APCI(+ve) 454/456 (M+H⁺).


m.p. 160-164°C.

^1H NMR (300 MHz, d_6 -DMSO) δ 8.25 (1H, t), 8.03 (1H, dd), 7.76 (1H, dd), 7.66 (1H, d), 7.52-7.41 (2H, m), 6.98 (1H, t), 4.69 (2H, s), 2.97 (2H, d), 1.94 (3H, s), 1.67 (3H, d), 1.59 (3H, d), 1.52 (6H, s).

5

Example 27

2-Chloro-5-[6-[(methylsulfonyl)amino]-2-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



- 10 **a) 2-Chloro-5-(6-chloro-2-pyridinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**
 Prepared according to the method of Example 14 (a) using [4-chloro-3-
 [[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a))
 (2.5 g) and 2,6-dichloropyridine (4.3 g) at room temperature. Purification (SiO_2 , 9:1
 isohexane:ethyl acetate as the eluant) gave the sub-title compound as a solid (1.1 g).

15

MS: APCI(+ve) 415/417 ($\text{M}+\text{H}^+$).

^1H NMR (300 MHz, CDCl_3) δ 8.29 (1H, d), 8.07 (1H, dd), 7.78 - 7.65 (2H, m), 7.51 (1H, d), 7.30 (1H, dd), 6.31 (1H, s), 3.21 (2H, d), 2.02 (3H, s), 1.80 - 1.59 (12H, m).

- 20 **b) 2-Chloro-5-[6-[(methylsulfonyl)amino]-2-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

2-Chloro-5-(6-chloro-2-pyridinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide
 (Example 27 (a)) (400 mg), methanesulfonamide (92 mg) and potassium carbonate (130

mg) were dissolved in methyl sulfoxide (3 mL). The mixture was heated to 190°C under nitrogen for 5 hours, then allowed to cool. Purification by chromatography (SiO₂, 3:1 *iso*hexane:ethyl acetate as eluant) and then by RP-HPLC (acetonitrile:aqueous ammonium acetate, Symmetry) afforded the title compound as a solid (19 mg).


MS: APCI(+ve) 474/476 (M+H⁺).

m.p. 115-120°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.43 (1H, t), 8.11-8.03 (2H, m), 7.80 (1H, t), 7.67-7.58 (2H, m), 6.88 (1H, d), 3.34 (3H, s), 2.97 (2H, d), 1.95 (3H, s), 1.68 (3H, d), 1.62 (3H, d), 1.55 (6H, s).

Example 28

[[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]oxy]-acetic acid



a) [[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]oxy]-acetic acid methyl ester

2-Chloro-5-(2-chloro-3-pyridinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 21 (a)) (100 mg), methyl glycolate (110 mg) and potassium *tert*-butoxide (1 mL, 1M solution in tetrahydrofuran) were placed in a 10 mL vial and heated at 70°C for 30 minutes in a microwave. The reaction mixture was allowed to cool and was concentrated.

Purification by chromatography (SiO₂, 3:1 *isohexane*:ethyl acetate as the eluant) afforded the sub-title compound as a solid (70 mg).

MS: APCI(+ve) 469/471(M+H⁺).

5

b) [[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]oxy]-acetic acid

[[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]oxy]-acetic acid methyl ester (Example 28 (a)) (70 mg) and methanol (1 mL) were placed in a 10 ml vial. A solution of potassium hydroxide (100 mg) in water (1 mL) was added and the mixture was heated at 50°C for 5 minutes in a microwave. The mixture was concentrated, water (10 mL) was added to the residue and this was then acidified to pH 2 with 2M aqueous hydrochloric acid. The resulting precipitate was collected by filtration. Purification (Varian NH₂ cartridge using dichloromethane (100 mL) and then 5 % trifluoroacetic acid in dichloromethane (100 mL) as eluant) afforded the title compound as a solid (17 mg).

10

15

MS: APCI(-ve) 453/455 (M-H⁺).

m.p. 226-229°C.


¹H NMR (400 MHz, d₆-DMSO) δ 8.40 (1H, t), 8.15 (1H, dd), 7.85 (1H, dd), 7.71 (1H, dd), 7.66 (1H, d), 7.57 (1H, d), 7.15 (1H, dd), 4.87 (2H, s), 2.95 (2H, d), 1.92 (3H, d), 1.67 (3H, d), 1.59 (3H, d), 1.53 (6H, s).

20

Example 29

2-Chloro-5-[3-(1*H*-tetrazol-5-ylmethoxy)-2-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

25



a) 2-Chloro-5-(3-hydroxy-2-pyridinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

Prepared according to the method of Example 14 (a) using [4-chloro-3-

5 [[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (2.0 g) and 2-bromo-3-hydroxypyridine (1.0 g) at room temperature to afford the sub-title compound as a solid (900 mg).

MS: APCI(+ve) 397/399 (M+H⁺).

10 ¹H NMR (400 MHz, d₆-DMSO) δ 10.38 (1H, s), 8.38 (1H, t), 8.17 (1H, dd), 8.14-8.09 (2H, m), 7.53 (1H, d), 7.36 (1H, dd), 7.24 (1H, dd), 2.96 (2H, d), 1.95 (3H, s), 1.68 (3H, d), 1.60 (3H, d), 1.54 (6H, s).

b) 2-Chloro-5-[3-(cyanomethoxy)-2-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

To a solution of 2-chloro-5-(3-hydroxy-2-pyridinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 29 (a)) (350 mg) and potassium carbonate (240 mg) in acetone (4 mL) was added chloroacetonitrile (0.1 mL). The mixture was heated at 55°C for 6 hours and was then allowed to cool and was concentrated. Purification by chromatography (SiO₂, 3:1 isohexane:ethyl acetate as eluant) afforded the sub-title compound as a solid (200 mg).

MS: APCI(+ve) 436/438 (M+H⁺).

c) 2-Chloro-5-[3-(1*H*-tetrazol-5-ylmethoxy)-2-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

Prepared according to the method of Example 9 using 2-chloro-5-[3-(cyanomethoxy)-2-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 29 (b)) (200 mg).

- 5 Purification (Varian NH₂ cartridge using 1:1 dichloromethane:methanol (100 mL) and then 1:5:5 acetic acid:dichloromethane:methanol (100 mL) as eluant) and further purification by RP-HPLC (acetonitrile:aqueous trifluoroacetic acid, Symmetry) afforded the title compound as a solid (69 mg).

10 MS: APCI(+ve) 479/481 (M+H⁺).


m.p. 122-126°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.38-8.30 (2H, m), 7.97 (1H, dd), 7.92 (1H, d), 7.74 (1H, d), 7.52 (1H, d), 7.44 (1H, dd), 5.63 (2H, s), 2.94 (2H, d), 1.93 (3H, s), 1.66 (3H, d), 1.58 (3H, d), 1.51 (6H, s).

15

Example 30

4'-Chloro-4-methoxy-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid



20

a) 2-Bromo-5-methoxy-benzoic acid ethyl ester

Prepared according to the method of Example 13 (a) using 2-bromo-5-methoxybenzoic acid (250 mg) and ethanol (2 mL) to afford the sub-title compound as a solid (280 mg).

MS: APCI(-ve) 258/260 (M-H⁺).

5

b) 4'-Chloro-4-methoxy-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid ethyl ester

Prepared according to the method of Example 14 (a) using [4-chloro-3-
[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a))
10 (250 mg) and 2-bromo-5-methoxy-benzoic acid ethyl ester (Example 30 (a)) (280 mg) at
room temperature to afford the sub-title compound as a solid (165 mg).

MS: APCI(+ve) 482/484 (M+H⁺).

¹H NMR (400 MHz, d₆-DMSO) δ 8.35 (1H, t), 7.50 (1H, d), 7.39 (1H, d), 7.32 (1H, dd),
15 7.27 (1H, d), 7.26-7.19 (2H, m), 4.09 (2H, q), 3.84 (3H, s), 2.94 (2H, d), 1.94 (3H, s), 1.67
(3H, d), 1.59 (3H, d), 1.52 (6H, s), 1.05 (3H, t).

c) 4'-Chloro-4-methoxy-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid

20 Prepared according to the method of Example 28 (b) using 4'-chloro-4-methoxy-3'-
[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid
ethyl ester (Example 30 (b)) (165 mg) to afford the title compound as a solid (75 mg).


MS: APCI(+ve) 454/456 (M+H⁺).

25 m.p. 219-221°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.35 (1H, t), 7.46 (1H, d), 7.38-7.27 (3H, m), 7.22 (1H,
d), 7.12 (1H, dd), 3.82 (3H, s), 2.94 (2H, d), 1.93 (3H, s), 1.66 (3H, d), 1.59 (3H, d), 1.52
(6H, s).

30 **Example 31**

4-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-1-methyl-1H-pyrazole-3-carboxylic acid



a) 2-Chloro-5-(3-formyl-1-methyl-1*H*-pyrazol-4-yl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

A mixture of [[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (300 mg), 4-bromo-1-methyl-1*H*-pyrazole-3-carboxaldehyde (160 mg), potassium carbonate (240 mg) and *tetrakis*(triphenylphosphine)palladium(0) (100 mg) in 1,2-dimethoxyethane (5 mL) / water (5 mL) was heated at 80°C with stirring under a nitrogen atmosphere for 3 hours. The products were filtered through diatomaceous earth, washing with methanol (3 x 10 mL), and the volatile components were removed *in vacuo*.

The residue was partitioned between dichloromethane (50 mL) and water (20 mL), the layers were separated and the aqueous fraction was extracted with dichloromethane (2 x 25 mL). The combined organic layers were washed with 2M aqueous hydrochloric acid (25 mL), saturated aqueous sodium hydrogen carbonate (25 mL) and saturated aqueous sodium chloride (25 mL) before being dried (MgSO₄), filtered and concentrated to give the subtitle compound as a solid (310 mg).

MS: APCI(+ve) 412/414 (M+H⁺).

b) 4-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-1-methyl-1*H*-pyrazole-3-carboxylic acid

To a solution of 2-chloro-5-(3-formyl-1-methyl-1*H*-pyrazol-4-yl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 31 (a)) (300 mg) in *N,N*-dimethylformamide (5 mL) was added potassium peroxydisulfate (670 mg) and the mixture was stirred at room temperature for 6 hours. The mixture was purified by Varian

NH₂ cartridge, eluting with methanol and then 5% trifluoroacetic acid in methanol. Further purification by RP-HPLC (acetonitrile:aqueous trifluoroacetic acid, Symmetry) gave the title compound as a colourless solid (50 mg).


5 MS: APCI(+ve) 428/430 (M+H⁺).

m.p. 166-175°C dec.

¹H NMR (400 MHz, d₆-DMSO) δ 8.32 (1H, t), 7.63 (1H, s), 7.57 (1H, dd), 7.52 (1H, d), 7.43 (1H, d), 3.99 (3H, s), 2.93 (2H, d), 1.94 (3H, s), 1.70-1.57 (6H, m), 1.54-1.50 (6H, m).

10 **Example 32**

4-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-1-methyl-1H-pyrazole-5-carboxylic acid



A mixture of [[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid
15 (Example 2 (a)) (350 mg), 4-bromo-1-methyl-1H-pyrazole-5-carboxaldehyde (190 mg),
potassium carbonate (300 mg) and *tetrakis*(triphenylphosphine)palladium(0) (50 mg) in
tetrahydrofuran (3 mL) / water (3 mL) was heated at 65°C with stirring under a nitrogen
atmosphere for 6 hours. The products were filtered through diatomaceous earth, washing
with methanol (3 x 10 mL) and the volatile components were removed *in vacuo*. The
20 residue was purified by chromatography (SiO₂, dichloromethane:methanol 99:1 as eluant)
and then redissolved in *N,N*-dimethylformamide (2 mL). Potassium peroxyxymonosulfate
(620 mg) was added, the mixture was stirred at room temperature for 18 hours and the
products were purified by Varian NH₂ cartridge, eluting with methanol and then 5%

trifluoroacetic acid in methanol. Further purification by recrystallisation from acetonitrile gave the title compound as a colourless solid (20 mg).


MS: APCI(+ve) 428 ($M+H^+$).

5 m.p. 187-190°C dec.

1H NMR (300 MHz, d_6 -DMSO) δ 8.35 (1H, t), 7.63 (1H, s), 7.51-7.41 (3H, m), 4.07 (3H, s), 2.93 (2H, d), 1.94 (3H, s), 1.72-1.55 (6H, m), 1.52 (6H, s).

Example 33

10 *N*-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]pyrazinyl]-*N*-methyl-glycine



a) *N*-(3-Chloropyrazinyl)-*N*-methyl-glycine, 1,1-dimethylethyl ester

15 A mixture of 2,3-dichloropyrazine (500 mg), *N*-methyl-glycine, 1,1-dimethylethyl ester hydrochloride (610 mg) and triethylamine (0.94 mL) in acetonitrile (1 mL) were heated in a microwave at 130°C for 3 hours. The mixture was then concentrated and purified by chromatography (SiO_2 , dichloromethane) to give the sub-title compound as an oil (400 mg).

20 MS: APCI(+ve) 202 ($M-C_4H_8+H^+$)

b) *N*-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]pyrazinyl]-*N*-methyl-glycine

A mixture of [[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (380 mg), *N*-(3-chloropyrazinyl)-*N*-methyl-glycine, 1,1-dimethylethyl ester (Example 33 (a)) (400 mg), potassium carbonate (300 mg) and *tetrakis*(triphenylphosphine)palladium(0) (60 mg) in tetrahydrofuran (3 mL) / water (3 mL) was stirred at room temperature under a nitrogen atmosphere for 16 hours. The products were filtered through diatomaceous earth, washing with methanol (3 x 10 mL) and the volatile components were removed *in vacuo*. The residue was partitioned between dichloromethane (25 mL) and water (10 mL), the layers were separated and the organic layer was dried (MgSO₄), filtered and concentrated. Purification by chromatography (SiO₂, *isohexane*:ethyl acetate 3:1, then *isohexane*:ethyl acetate 1:1) gave an oil which was dissolved in dichloromethane (5 mL). Trifluoroacetic acid (2 mL) was added, the mixture was stirred at room temperature for 24 hours and then concentrated. Purification by Varian NH₂ resin, eluting with methanol and then 5% trifluoroacetic acid in methanol, and then by trituration with diethyl ether, gave the title compound as a solid (13 mg).


MS: APCI(+ve) 469 (M+H⁺).

m.p. 151-153°C dec.

¹H NMR (400 MHz, d₆-DMSO) δ 8.38 (1H, t), 8.12 (1H, d), 8.09 (1H, d), 7.77 (1H, dd), 7.71 (1H, d), 7.57 (1H, d), 4.08 (2H, s), 2.95 (2H, d), 2.70 (s, 3H), 1.94 (3H, s), 1.71-1.56 (6H, m), 1.55-1.50 (6H, m).

Example 34

1-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]pyrazinyl]-4-piperidinecarboxylic acid



a) 1-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-

ylmethyl)amino]carbonyl]phenyl]pyrazinyl]-4-piperidinecarboxylic acid, ethyl ester

A mixture of 2-chloro-5-(3-chloropyrazinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 5 benzamide (Example 5 (a)) (200 mg), ethyl isonipecotatate (230 mg) and acetonitrile (1 mL)
 was heated in a microwave at 140°C for 6 hours. The products were partitioned between
 dichloromethane (25 mL) and 2M aqueous hydrochloric acid (10 mL), the layers were
 separated and the aqueous layer was extracted with dichloromethane (25 mL). The
 combined organics were washed with saturated aqueous sodium chloride (10 mL), dried
 10 (MgSO₄), filtered and concentrated *in vacuo*. Purification by chromatography (SiO₂,
 dichloromethane:methanol 99.5:0.5 as eluant) gave the sub-title compound as a film (140
 mg).

MS: APCI(+ve) 537 (M+H⁺)

b) 1-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-

ylmethyl)amino]carbonyl]phenyl]pyrazinyl]- 4-piperidinecarboxylic acid

Prepared according to the method of Example 23 (d) using 1-[3-[4-chloro-3-
 [[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]pyrazinyl]-4-
 20 piperidinecarboxylic acid, ethyl ester (Example 34 (a)) (140 mg). Purification by
 chromatography (SiO₂, dichloromethane:methanol 98:2 as eluant) gave the title compound
 as a solid (86 mg).

MS: APCI(+ve) 509 (M+H⁺).


m.p. 136-140°C.

¹H NMR (300 MHz, d₆-DMSO) δ 12.22 (1H, s), 8.39 (1H, t), 8.22 (1H, d), 8.20 (1H, d), 8.00 (1H, dd), 7.95 (1H, d), 7.60 (1H, d), 3.56-3.45 (2H, m), 2.95 (2H, d), 2.83-2.71 (2H, m), 2.44-2.31 (1H, m), 1.99-1.90 (3H, m), 1.85-1.75 (2H, m), 1.72-1.47 (14H, m).

5

Example 35

4'-Chloro-6-fluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid



10 a) 2-Bromo-3-fluoro-benzoic acid, methyl ester

Prepared according to the method of Example 13 (a) using 2-bromo-3-fluoro-benzoic acid (500 mg). Purification by chromatography (SiO₂, dichloromethane as eluant) gave the subtitle compound as a colourless oil (480 mg).

15 ¹H NMR (400 MHz, CDCl₃) δ 7.58 (1H, ddd), 7.34 (1H, ddd), 7.26 (1H, ddd), 3.95 (3H, s).

b) 4'-Chloro-6-fluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid, methyl ester

20 A mixture of [[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (500 mg), 2-bromo-3-fluoro-benzoic acid, methyl ester (Example 35 (a)) (340 mg), potassium carbonate (400 mg) and *tetrakis*(triphenylphosphine)palladium(0) (80 mg) in tetrahydrofuran (3 mL) / water (3 mL) was stirred at 65°C under a nitrogen atmosphere for 16 hours. The products were filtered through diatomaceous earth, washing

with methanol (3 x 10 mL) and the volatile components were removed *in vacuo*. The residue was partitioned between dichloromethane (20 mL) and 2M aqueous hydrochloric acid (10 mL), the layers were separated, the aqueous fraction was extracted with dichloromethane (2 x 20 mL) and the combined organic layers were washed with saturated aqueous sodium chloride (10 mL) before being dried (MgSO₄), filtered and concentrated. Purification by chromatography (SiO₂, dichloromethane) gave the sub-title compound as a colourless solid (250 mg).

MS: APCI(+ve) 456 (M+H⁺).

c) 4'-Chloro-6-fluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid

A solution of sodium hydroxide (66 mg) in water (0.5 mL) was added to a solution of 4'-chloro-6-fluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid, methyl ester (Example 35 (b)) (250 mg) in tetrahydrofuran (2 mL) in a 10 mL microwave vial and this was heated at 70°C for 3 hours within a microwave. The reaction mixture was concentrated and then acidified to pH 5 using 2M aqueous hydrochloric acid. Purification by chromatography (SiO₂, dichloromethane, then dichloromethane:methanol 98:2 as eluant) gave the title compound as a colourless solid (55 mg).


MS: APCI(-ve) 440 (M-H⁺).

m.p. 108-113°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.39 (1H, t), 7.68-7.62 (1H, m), 7.59-7.46 (3H, m), 7.37 (1H, d), 7.29 (1H, d), 2.93 (2H, d), 1.93 (3H, s), 1.70-1.55 (6H, m), 1.52 (6H, s).

Example 36

4'-Chloro-5-fluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid



To a stirred solution of 2-bromo-4-fluoro-benzoic acid (500 mg) in dichloromethane (5 mL) at 0°C under nitrogen was added *N,N*-dimethylformamide (1 drop) and oxalyl chloride (0.4 mL). The reaction mixture was allowed to warm to room temperature over 2 hours, concentrated *in vacuo* and then redissolved in dichloromethane (5 mL). Methanol (1 mL) was added dropwise and the mixture was stirred at room temperature for 2 hours before being concentrated *in vacuo* to give a colourless oil which was dissolved in tetrahydrofuran (3 mL) / water (3 mL). [[(Tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (500 mg), potassium carbonate (400 mg) and *tetrakis*(triphenylphosphine)palladium(0) (80 mg) were added and the mixture was stirred at 65°C under a nitrogen atmosphere for 8 hours. The products were filtered through diatomaceous earth, washing with methanol (3 x 10 mL), and were then concentrated *in vacuo*. Dichloromethane (20 mL) and 2M aqueous hydrochloric acid (10 mL) were added, the layers were separated and the aqueous fraction was extracted with dichloromethane (2 x 10 mL). The combined organic layers were washed with saturated aqueous sodium chloride (10 mL), dried (MgSO₄), filtered and concentrated. The residue was dissolved in tetrahydrofuran (3 mL) and a solution of sodium hydroxide (250 mg) in water (3 mL) was added. The mixture was stirred at room temperature for 24 hours, adjusted to pH 5 with 2M aqueous hydrochloric acid and concentrated *in vacuo*. Purification by chromatography (SiO₂, dichloromethane:methanol 98:2 as eluant) gave the title compound as a colourless solid (90 mg).


MS: APCI(-ve) 440 (M-H⁺).

m.p. 120-124°C.

^1H NMR (400 MHz, d_6 -DMSO) δ 13.01 (1H, s), 8.36 (1H, t), 7.87 (1H, dd), 7.52 (1H, d), 7.42-7.37 (2H, m), 7.34 (1H, ddd), 7.27 (1H, dd), 2.95 (2H, d), 1.94 (3H, s), 1.71-1.56 (6H, m), 1.55-1.50 (6H, m).

5 **Example 37**

4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-acetic acid



a) 2-Bromo-benzeneacetic acid, methyl ester

10 Prepared according to the method of Example 13 (a) using 2-bromo-benzeneacetic acid (300 mg), oxalyl chloride (0.4 mL), *N,N*-dimethylformamide (1 drop), dichloromethane (2 mL) and methanol (2 mL), to give the sub-title compound as a solid (315 mg).

MS: APCI(+ve) 230 ($\text{M}+\text{H}^+$).

15

b) 4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-acetic acid, methyl ester

A mixture of [4-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (250 mg), 2-bromo-benzeneacetic acid, methyl ester (Example 37 (a)) (198 mg), potassium carbonate (199 mg) and
 20 bis(triphenylphosphine)palladium(II) chloride (50 mg) in tetrahydrofuran (2 mL) / water (1 mL) was heated at 40°C under a nitrogen atmosphere for 1 hour. The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO_2 , dichloromethane as eluant) to yield the sub-title compound as a solid (100 mg).

MS: APCI(+ve) 452/454 ($M+H^+$).

c) 4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-acetic acid

Prepared according to the method of Example 10 (b) using 4'-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-acetic acid, methyl ester (Example 37 (b)) (100 mg), potassium hydroxide (100 mg), methanol (1 mL) and water (1 mL). Purification (Varian NH₂ cartridge using methanol (100 mL) and then 5 % acetic acid in methanol (100 mL) as eluant) afforded the title compound as a solid (19 mg).


MS: APCI(+ve) 438/440 ($M+H^+$).

m.p. 200-202°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.36 (1H, t), 7.55 (1H, d), 7.39-7.32 (4H, m), 7.31 (1H, d), 7.27-7.23 (1H, m), 3.52 (2H, s), 2.94 (2H, d), 1.93 (3H, s), 1.66 (3H, d), 1.59 (3H, d), 1.52 (6H, s).

Example 38

[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-3-yl]oxy]-acetic acid



a) 4-Chloro-3'-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-[1,1'-biphenyl]-3-carboxamide

Prepared according to the method of Example 37 (b) using [4-chloro-3-
[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a))
(1 g), 3-bromo-phenol (600 mg), potassium carbonate (800 mg),
bis(triphenylphosphine)palladium(II) chloride (200 mg), tetrahydrofuran (10 mL) and
5 water (10 mL). Purification by chromatography (SiO₂, 1:3 ethyl acetate:*iso*hexane as
eluant) gave the sub-title compound as a solid (650 mg).

MS: APCI(+ve) 396/398 (M+H⁺).

m.p. 182-184°C.

10 ¹H NMR (300 MHz, d₆-DMSO) δ 9.59 (1H, s), 8.40 (1H, t), 7.64 (1H, dd), 7.58-7.49 (2H,
m), 7.30 (1H, t), 7.09 (1H, d), 7.03 (1H, s), 6.80 (1H, dd), 2.96 (2H, d), 1.95 (3H, s), 1.76-
1.46 (12H, m).

15 **b) [[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-
3-yl]oxy]-acetic acid, ethyl ester**

Prepared according to the method of Example 15 (a) using 4-chloro-3'-hydroxy-*N*-
(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-[1,1'-biphenyl]-3-carboxamide (Example 38 (a)) (250
mg), ethyl chloroacetate (300 mg), potassium carbonate (174 mg) and acetone (4 mL).
Purification by chromatography (SiO₂, dichloromethane as eluant) gave the sub-title
20 compound as a solid (180 mg).

MS: APCI(+ve) 482/484 (M+H⁺).

25 ¹H NMR (300 MHz, d₆-DMSO) δ 8.39 (1H, t), 7.72 (1H, dd), 7.65 (1H, d), 7.56 (1H, d),
7.40 (1H, t), 7.29 (1H, d), 7.23 (1H, t), 6.97 (1H, dd), 4.88 (2H, s), 4.18 (2H, q), 2.97 (2H,
d), 1.95 (3H, s), 1.68 (3H, d), 1.60 (3H, d), 1.54 (6H, s), 1.21 (3H, t).

**c) [[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-
3-yl]oxy]-acetic acid**

Prepared according to the method of Example 15 (b) using [[4'-Chloro-3'-
30 [[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-3-yl]oxy]-acetic acid,
ethyl ester (Example 38 (b)) (180 mg), potassium hydroxide (100 mg), methanol (1 mL)

and water (1 mL). Purification by RP-HPLC (acetonitrile:aqueous trifluoroacetic acid, Symmetry) gave the title compound as a solid (22 mg).


MS: APCI(+ve) 454/456 ($M+H^+$).

5 m.p. 99-101°C.

1H NMR (400 MHz, d_6 -DMSO) δ 8.40 (1H, t), 7.72 (1H, dd), 7.65 (1H, d), 7.56 (1H, d), 7.40 (1H, t), 7.27 (1H, d), 7.21 (1H, t), 6.95 (1H, dd), 4.78 (2H, s), 2.97 (2H, d), 1.95 (3H, s), 1.68 (3H, d), 1.60 (3H, d), 1.54 (6H, s).

10 Example 39

(2*R*)-2-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid



15 **a) (2*R*)-2-[[4'-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid, methyl ester**

Prepared according to the method of Example 15 (a) using 4-chloro-2'-hydroxy-*N*-(tricyclo[3.3.1.1^{3,7}]]dec-1-ylmethyl)-[1,1'-biphenyl]-3-carboxamide (Example 14 (a)) (170 mg), methyl (2*S*)-2-chloropropanoate (212 mg), potassium carbonate (120 mg) and acetone (4 mL). Purification by chromatography (SiO_2 , 1:4 ethyl acetate:*iso*hexane) gave the sub-
20 title compound as a solid (140 mg).

MS: APCI(+ve) 482/484 ($M+H^+$).

¹H NMR (400 MHz, d₆-DMSO) δ 8.39 (1H, t), 7.67-7.61 (2H, m), 7.52 (1H, d), 7.40-7.28 (2H, m), 7.07 (1H, t), 6.96 (1H, d), 5.10 (1H, q), 3.67 (3H, s), 2.99-2.90 (2H, m), 1.93 (3H, s), 1.67 (3H, d), 1.59 (3H, d), 1.53 (6H, s), 1.45 (3H, d).

5 **b) (2*R*)-2-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid**

Prepared according to the method of Example 10 (b) using (2*R*)-2-[[4'-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid, methyl ester (Example 39 (a)) (140 mg), potassium hydroxide (100 mg), methanol (2
10 mL) and water (1 mL). Purification (Varian NH₂ cartridge using methanol (100 mL) and then 5 % acetic acid in methanol (100 mL) as eluant) afforded the title compound as a solid (100 mg).


MS: APCI(+ve) 468/470 (M+H⁺).

15 m.p. 133-137°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.43 (1H, t), 7.75 (1H, dd), 7.67 (1H, d), 7.48 (1H, d), 7.31 (1H, d), 7.25 (1H, t), 6.97 (1H, t), 6.90 (1H, d), 4.64 (1H, d), 2.94 (2H, d), 1.92 (3H, s), 1.66 (3H, d), 1.59 (3H, d), 1.52 (6H, s), 1.35 (3H, d).

20 **Example 40**

[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-4-yl]oxy]-acetic acid



a) 4-chloro-4'-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-[1,1'-biphenyl]-3-carboxamide

A mixture of [4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (1.0 g), 4-bromo-phenol (600 mg), potassium carbonate (800 mg) and *bis*(triphenylphosphine)palladium(II) chloride (200 mg) in tetrahydrofuran (10 mL) / water (10 mL) was stirred at room temperature under a nitrogen atmosphere for 16 hours. The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO₂, 1:3 ethyl acetate:*isohexane*) to yield the sub-title compound as a solid (470 mg).

MS: APCI(+ve) 396/398 (M+H⁺).

b) [[4'-chloro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl][1,1'-biphenyl]-4-yl]oxy]-acetic acid, ethyl ester

Prepared according to the method of Example 15 (a) using 4-chloro-4'-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-[1,1'-biphenyl]-3-carboxamide (Example 40 (a)) (250 mg), ethyl chloroacetate (300 mg), potassium carbonate (174 mg) and acetone (4 mL). Purification by chromatography (SiO₂, 1:4 ethyl acetate:*isohexane* as eluant) gave the sub-title compound as a solid (190 mg).

MS: APCI(+ve) 482/484 (M+H⁺).

c) **[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-4-yl]oxy]-acetic acid**

Prepared according to the method of Example 10 (b) using [[4'-chloro-3'-
[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-4-yl]oxy]-acetic acid,
ethyl ester (Example 40 (b)) (190 mg), potassium hydroxide (100 mg), methanol (2 mL)
and water (1 mL). Purification by RP-HPLC (acetonitrile:aqueous ammonium acetate,
Symmetry) afforded the title compound as a solid (74 mg).


MS: APCI(+ve) 454/456 (M+H⁺).

m.p. 136-140°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.38 (1H, t), 7.64 (1H, dd), 7.60-7.55 (3H, m), 7.50 (1H, d), 6.92 (2H, d), 4.34 (2H, s), 2.96 (2H, d), 1.95 (3H, s), 1.67 (3H, d), 1.60 (3H, d), 1.54 (6H, s).

Example 41

(2S)-2-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-3-yl]oxy]-propanoic acid



a) **(2S)-2-[[4'-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-3-yl]oxy]- propanoic acid, methyl ester**

Prepared according to the method of Example 15 (a) using 4-chloro-3'-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-[1,1'-biphenyl]-3-carboxamide (Example 38 (a)) (250 mg), methyl (2R)-2-chloropropanoate (311 mg), potassium carbonate (174 mg) and

acetone (4 mL). Purification by chromatography (SiO₂, 1:4 ethyl acetate:isohexane) gave the sub-title compound as a solid (200 mg).

MS: APCI(+ve) 482/484 (M+H⁺).

5 ¹H NMR (400 MHz, d₆-DMSO) δ 8.39 (1H, t), 7.70 (1H, dd), 7.63 (1H, d), 7.56 (1H, d), 7.39 (1H, t), 7.28 (1H, d), 7.20 (1H, s), 6.92 (1H, dd), 5.16 (1H, q), 3.68 (3H, s), 2.97 (2H, d), 1.95 (3H, s), 1.73 - 1.48 (15H, m).

10 **b) (2*S*)-2-[[4'-Chloro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-3-yl]oxy]-propanoic acid**

Prepared according to the method of Example 10 (b) using (2*S*)-2-[[4'-chloro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-3-yl]oxy]-propanoic acid, methyl ester (Example 41 (a)) (200 mg), potassium hydroxide (100 mg), methanol (2 mL) and water (1 mL) to afford the title compound as a solid (175 mg).


15 MS: APCI(+ve) 468/470 (M+H⁺).

m.p. 115-120°C.

20 ¹H NMR (400 MHz, d₆-DMSO) δ 8.39 (1H, t), 7.69 (1H, dd), 7.62 (1H, d), 7.54 (1H, d), 7.37 (1H, t), 7.24 (1H, d), 7.16 (1H, s), 6.89 (1H, dd), 4.93 (1H, q), 2.96 (2H, d), 1.93 (3H, d), 1.73-1.43 (15H, m).

Example 42

4,4'-Dichloro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid



a) 4,4'-Dichloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid, methyl ester

Prepared according to the method of Example 40 (a) using [4-chloro-3-

5 [[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (250 mg), methyl 2-bromo-5-chlorobenzoate (215 mg), potassium carbonate (200 mg), *bis*(triphenylphosphine)palladium(II) chloride (50 mg), tetrahydrofuran (2 mL) and water (2 mL). Purification by chromatography (SiO₂, 1:4 ethyl acetate:*iso*hexane) gave the subtitle compound as a solid (200 mg).

10 MS: APCI(+ve) 472/474 (M+H⁺).

¹H NMR (400 MHz, d₆-DMSO) δ 8.40 (1H, t), 7.83 (1H, d), 7.74 (1H, dd), 7.58-7.47 (2H, m), 7.36 (1H, dd), 7.30 (1H, d), 3.65 (3H, d), 2.95 (2H, d), 1.94 (3H, s), 1.67 (3H, d), 1.60 (3H, d), 1.52 (6H, s).

b) 4,4'-Dichloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid

Prepared according to the method of Example 10 (b) using 4,4'-dichloro-3'-

20 [[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid, methyl ester (Example 42 (a)) (200 mg), potassium hydroxide (100 mg), methanol (2 mL) and water (1 mL). Purification (Varian NH₂ cartridge using methanol (100 mL) and then 5 % acetic acid in methanol (100 mL) as eluant) afforded the title compound as a solid (125 mg).


MS: APCI(-ve) 456 (M-H⁺).

m.p. 207-210°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.38 (1H, t), 7.78 (1H, d), 7.68 (1H, dd), 7.52 (1H, d),
5 7.44 (1H, d), 7.40-7.33 (2H, m), 2.94 (2H, d), 1.94 (3H, s), 1.67 (3H, d), 1.59 (3H, d), 1.52
(6H, s).

Example 43

(2*S*)-2-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-
10 biphenyl]-4-yl]oxy]-propanoic acid



a) (2*S*)-2-[[4'-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-
biphenyl]-4-yl]oxy]-propanoic acid, methyl ester

Prepared according to the method of Example 15 (a) using 4-chloro-4'-hydroxy-*N*-
15 (tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-[1,1'-biphenyl]-3-carboxamide (Example 40 (a)) (210
mg), methyl (2*R*)-2-chloropropanoate (262 mg), potassium carbonate (150 mg) and
acetone (4 mL). Purification by chromatography (SiO₂, 1:2 ethyl acetate:*isohexane*) gave
the sub-title compound as a solid (140 mg).

20 MS: APCI(+ve) 482/484 (M+H⁺).

b) (2*S*)-2-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-4-yl]oxy]-propanoic acid

Prepared according to the method of Example 10 (b) using (2*S*)-2-[[4'-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-4-yl]oxy]-propanoic acid, methyl ester (Example 43 (a)) (140 mg), potassium hydroxide (100 mg), methanol (2 mL) and water (1 mL). Purification by RP-HPLC (acetonitrile:aqueous ammonium acetate, Symmetry) afforded the title compound as a solid (50 mg).


MS: APCI(+ve) 468/470 (M+H⁺).

m.p. 145-150°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.38 (1H, t), 7.63 (1H, dd), 7.60-7.53 (3H, m), 7.50 (1H, d), 6.91 (2H, d), 4.59 (1H, q), 2.96 (2H, d), 1.94 (3H, s), 1.67 (3H, d), 1.60 (3H, d), 1.54 (6H, s), 1.43 (3H, d).

Example 44

3-Chloro-6-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid



a) 3-Chloro-6-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-

ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid, methyl ester

Prepared according to the method of Example 40 (a) using [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (200 mg), methyl 3,6-dichloropyridine-2-carboxylate (123 mg), potassium carbonate (166

mg), *bis*(triphenylphosphine)palladium(II) chloride (42 mg), tetrahydrofuran (2 mL) and water (2 mL). Purification by chromatography (SiO₂, 1:4 ethyl acetate:*isohexane*) gave the sub-title compound as a solid (130 mg).

5 MS: APCI(+ve) 473/475 (M+H⁺).

b) 3-Chloro-6-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid

Prepared according to the method of Example 10 (b) using 3-chloro-6-[4-chloro-3-
10 [[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid, methyl ester (Example 44 (a)) (130 mg), potassium hydroxide (100 mg), methanol (2 mL) and water (1 mL). Purification by chromatography (SiO₂, 1:9 methanol:dichloromethane) gave the title compound as a solid (100 mg).


15 MS: APCI(+ve) 459/461 (M+H⁺).

m.p. 154-159°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.45 (1H, t), 8.24-8.09 (4H, m), 7.64 (1H, d), 2.98 (2H, d), 1.95 (3H, s), 1.68 (3H, d), 1.61 (3H, d), 1.55 (6H, s).

20 **Example 45**

3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-pyridinecarboxylic acid



a) 3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-pyridinecarboxylic acid, methyl ester

Prepared according to the method of Example 37 (b) using [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (250 mg), 3-iodo-4-pyridinecarboxylic acid, methyl ester (190 mg), potassium carbonate (200 mg), *bis*(triphenylphosphine)palladium(II) chloride (51 mg), tetrahydrofuran (2 mL) and water (2 mL). Purification by chromatography (SiO₂, 1:3 ethyl acetate:*iso*hexane as eluant) gave the sub-title compound as a solid (80 mg).

MS: APCI(+ve) 439/441 (M+H⁺).

¹H NMR (300 MHz, CDCl₃) δ 8.75 (1H, d), 8.67 (1H, d), 7.72-7.66 (2H, m), 7.48 (1H, d), 7.33 (1H, dd), 6.38-6.27 (1H, m), 3.76 (3H, s), 3.19 (2H, d), 2.02 (3H, s), 1.79 - 1.54 (12H, m).

b) 3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-pyridinecarboxylic acid

Prepared according to the method of Example 10 (b) using 3-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-pyridinecarboxylic acid, methyl ester (Example 45 (a)) (80 mg), potassium hydroxide (100 mg), methanol (2 mL) and water (1 mL). Purification by chromatography (SiO₂, 1:9 methanol:dichloromethane as eluant) gave the title compound as a solid (30 mg).


MS: APCI(-ve) 423/425 (M-H⁺).

m.p. 170-240°C dec.

¹H NMR (400 MHz, d₆-DMSO) δ 8.64-8.55 (2H, m), 8.38 (1H, t), 7.56-7.44 (4H, m), 2.94 (2H, d), 1.93 (3H, s), 1.66 (3H, d), 1.59 (3H, d), 1.53 (6H, s).

Example 46

[[2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinyl]oxy]-acetic acid



a) 2-Chloro-5-(3-hydroxy-2-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

Prepared according to the method of Example 26 (a) using [4-chloro-3-

- 5 [[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (2.0 g), 3-hydroxy-2-bromopyridine (1.0 g), sodium carbonate (1.27 g), *tetrakis*(triphenylphosphine)palladium(0) (693 mg), toluene (20 mL), ethanol (20 mL) and water (20 mL). Purification by chromatography (SiO₂, 1:1 ethyl acetate:*isohexane* as eluant) gave the sub-title compound as a solid (900 mg).

10 MS: APCI(+ve) 397/399 (M+H⁺).

¹H NMR (400 MHz, d₆-DMSO) δ 10.38 (1H, s), 8.38 (1H, t), 8.17 (1H, dd), 8.14 – 8.09 (2H, m), 7.53 (1H, d), 7.36 (1H, dd), 7.24 (1H, dd), 2.96 (2H, d), 1.95 (3H, s), 1.68 (3H, d), 1.60 (3H, d), 1.54 (6H, s).

b) [[2-[4-Chloro-3-[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinyl]oxy]-acetic acid, ethyl ester

Prepared according to the method of Example 15 (a) using 2-chloro-5-(3-hydroxy-2-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 46 (a)) (250 mg), ethyl chloroacetate (300 mg), potassium carbonate (174 mg) and acetone (4 mL).

20 Purification by chromatography (SiO₂, 1:2 ethyl acetate:*isohexane* as eluant) gave the sub-title compound as a solid (150 mg).

MS: APCI(+ve) 483/485 (M+H⁺).

¹H NMR (400 MHz, d₆-DMSO) δ 8.38 (1H, t), 8.31 (1H, dd), 8.07 (1H, dd), 8.03 (1H, d), 7.57-7.52 (2H, m), 7.38 (1H, dd), 4.95 (2H, s), 4.18 (2H, q), 2.95 (2H, d), 1.94 (3H, s), 1.67 (3H, d), 1.60 (3H, d), 1.53 (6H, s), 1.21 (3H, t).

5 **c) [[2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinyl]oxy]-acetic acid**

Prepared according to the method of Example 10 (b) using [[2-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinyl]oxy]-acetic acid, ethyl ester (Example 46 (b)) (150 mg), potassium hydroxide (100 mg), methanol (2 mL)
10 and water (1 mL). Purification (Varian NH₂ cartridge using dichloromethane (100 mL) and then 5 % acetic acid in dichloromethane (100 mL) as eluant) afforded the title compound as a solid (139 mg).


MS: APCI(-ve) 453/455 (M-H⁺).

15 m.p. 133-137°C.

¹H NMR (300 MHz, d₆-DMSO) δ 8.38 (1H, t), 8.28 (1H, dd), 8.07 (1H, dd), 8.02 (1H, d), 7.54 (1H, d), 7.49 (1H, dd), 7.36 (1H, dd), 4.80 (2H, s), 2.95 (2H, d), 1.94 (3H, s), 1.67 (3H, d), 1.59 (3H, d), 1.53 (6H, s).

20 **Example 47**

N-[2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinyl]-glycine



a) 2-Chloro-5-(3-nitro-2-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

Prepared according to the method of Example 37 (b) using [4-chloro-3-
[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a))
(1.0 g), 2-chloro-3-nitropyridine (460 mg), potassium carbonate (800 mg),
bis(triphenylphosphine)palladium(II) chloride (100 mg), tetrahydrofuran (10 mL) and
5 water (10 mL). Purification by chromatography (SiO₂, 1:9 ethyl acetate:isohexane as
eluant) gave the sub-title compound as a solid (600 mg).

MS: APCI(+ve) 426/428 (M+H⁺).

¹H NMR (400 MHz, d₆-DMSO) δ 8.95 (1H, dd), 8.54-8.44 (2H, m), 7.76-7.71 (1H, m),
10 7.64-7.55 (3H, m), 2.96 (2H, t), 1.94 (3H, s), 1.67 (3H, d), 1.60 (3H, d), 1.53 (6H, s).

b) 5-(3-Amino-2-pyridinyl)-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

Iron powder (600 mg) was added portionwise to a stirred mixture of 2-chloro-5-(3-nitro-2-
pyridinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 47 (a)) (600 mg),
15 ammonium chloride (600 mg), ethanol (5 mL) and water (5 mL). The mixture was then
heated at 50 °C for 24 hours then allowed to cool and filtered through diatomaceous earth,
washing with dichloromethane (200 mL). The filtrate and washings were evaporated and
the combined residues were partitioned between dichloromethane (100 mL) and water (100
mL). The layers were separated and the organics were dried, filtered and evaporated to
20 afford the sub-title compound as a solid (450 mg).

MS: APCI(+ve) 396/398 (M+H⁺).

¹H NMR (400 MHz, d₆-DMSO) δ 8.37 - 8.26 (1H, m), 7.92 (1H, dd), 7.78 - 7.67 (2H, m),
7.59 - 7.52 (1H, m), 7.15 (1H, dd), 7.09 (1H, dd), 5.26 (2H, s), 2.96 (2H, d), 1.95 (3H, s),
25 1.74 - 1.46 (12H, m).

c) *N*-[2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinyl]-glycine

Ethyl glyoxalate (50 % solution in toluene) (200 mg) was added to a stirred mixture of 5-
30 (3-amino-2-pyridinyl)-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide
(Example 47 (b)) (400 mg) and activated 3 Å molecular sieves (500 mg) in
dichloromethane (5 mL). The mixture was stirred at room temperature under nitrogen for 3

hours, sodium triacetoxyborohydride (642 mg) was added, stirring was continued for a further 16 hours and the mixture was then concentrated *in vacuo*. Purification by chromatography (SiO₂, ethyl acetate, then 1:9 methanol:ethyl acetate as eluant) afforded the title compound (18 mg).

5

MS: APCI(-ve) 452/454 (M-H⁺).


m.p. 190-194°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.47 (1H, t), 7.88 (1H, d), 7.70 (1H, dd), 7.64 (1H, d), 7.58 (1H, d), 7.15 (1H, d), 6.90 (1H, d), 5.49 (1H, s), 3.48-3.16 (2H, m), 2.95 (2H, d), 1.93 (3H, s), 1.66 (3H, d), 1.59 (3H, d), 1.52 (6H, s).

10

Example 48

4'-Chloro-4,5-difluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid



15

a) 2-Bromo-4,5-difluoro-benzoic acid, methyl ester

Prepared according to the method of Example 13 (a) using 2-bromo-4,5-difluorobenzoic acid (237 mg), oxalyl chloride (0.1 mL), *N,N*-dimethylformamide (1 drop), dichloromethane (2 mL) and methanol (2 mL) to give the sub-title compound as a solid (250 mg).

20

¹H NMR (300 MHz, CDCl₃) δ 7.74 (1H, dd), 7.51 (1H, dd), 3.93 (3H, s).

b) 4'-Chloro-4,5-difluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid, methyl ester

Prepared according to the method of Example 40 (a) using [4-chloro-3-
[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a))
5 (200 mg), 2-bromo-4,5-difluoro-benzoic acid, methyl ester (Example 48 (a)) (175 mg),
potassium carbonate (170 mg), *bis*(triphenylphosphine)palladium(II) chloride (20 mg),
tetrahydrofuran (1 mL) and water (1 mL). Purification by chromatography (SiO₂, 1:4 ethyl
acetate:isohexane as eluant) gave the sub-title compound as a solid (60 mg).

10 MS: APCI(+ve) 474 (M+H⁺).

c) 4'-Chloro-4,5-difluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid

A solution of potassium hydroxide (50 mg) in water (0.5 mL) was added to a solution of
15 4'-chloro-4,5-difluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-
biphenyl]-2-carboxylic acid, methyl ester (Example 48 (b)) (60 mg), in methanol (0.5 mL)
and tetrahydrofuran (0.5 mL). The mixture was stirred at room temperature for 2 hours
then concentrated. The residue was dissolved in water (5 mL) and the solution was
acidified to pH 2 with 2 M aqueous hydrochloric acid. The resulting solid was collected by
20 filtration and washed with water to afford the title compound (50 mg).


MS: APCI(-ve) 458 (M-H⁺).

m.p. 121-125°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.35 (1H, t), 7.84 (1H, dd), 7.56-.48 (2H, m), 7.41-7.36
25 (2H, m), 2.94 (2H, d), 1.94 (3H, s), 1.67 (3H, d), 1.59 (3H, d), 1.52 (6H, s).

Example 49

4'-Chloro-3'-[[[(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid



a) 2-Chloro-5-iodo-benzoic acid, 1,1-dimethylethyl ester

N,N-Dimethylformamide (1 drop) and oxalyl chloride (4.8 mL) were added to a stirred solution of 2-chloro-5-iodobenzoic acid (5 g) in dichloromethane (20 mL) at 0 °C. The reaction was allowed to warm to room temperature, stirred under nitrogen for 2 hours, and then evaporated to dryness. The residue was dissolved in tetrahydrofuran (20 mL) and cooled to 0 °C. Potassium *tert*-butoxide (22 mL, 1 M solution in tetrahydrofuran) was added over 10 minutes. The reaction was allowed to warm to room temperature and stirred under nitrogen for 2 hours then poured into saturated aqueous sodium bicarbonate (50 mL). The layers were separated and the aqueous was extracted with diethyl ether (50 mL). The combined organics were dried, filtered and evaporated to afford the sub-title compound as an oil (5.7 g).

¹H NMR (400 MHz, d₆-DMSO) δ 7.99 (1H, d), 7.87 (1H, dd), 7.34 (1H, d), 1.54 (9H, s).

b) 2-Chloro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-benzoic acid, 1,1-dimethylethyl ester

A mixture of 2-Chloro-5-iodo-benzoic acid, 1,1-dimethylethyl ester (Example 49 (a)) (5 g), *bis*(pinacolato)diboron (6 g), [1,1'-*bis*(diphenylphosphino)ferrocene]dichloropalladium(II)dichloromethane (600 mg) and potassium acetate (6.5 g) in *N,N*-dimethylformamide (50 mL) was heated to 90 °C under nitrogen for 90 minutes. The mixture was allowed to cool then diluted with 2:1 ethyl acetate: diethyl ether (250 mL) and filtered through diatomaceous earth. The filtrate was

washed with water (250 mL) and brine (100 mL) then evaporated. Purification (SiO₂, 1:1 diethyl ether : *isohexane* as eluant) afforded the sub-title compound as a solid (5.5 g).

MS: APCI(+ve) 282 (M-C₄H₈+H⁺).

5 ¹H NMR (300 MHz, d₆-DMSO) δ 7.88 (1H, d), 7.76 (1H, dd), 7.56 (1H, d), 1.55 (9H, s), 1.32 (12H, s).

c) 4'-Chloro-[1,1'-biphenyl]-2,3'-dicarboxylic acid, 3'-(1,1-dimethylethyl) 2-methyl ester

10 Prepared according to the method of Example 40 (a) using 2-chloro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-benzoic acid, 1,1-dimethylethyl ester (Example 49 (b)) (3.5 g), methyl-2-bromobenzoate (2.23 g), potassium carbonate (2.87 g), *bis*(triphenylphosphine)palladium(II) chloride (365 mg), tetrahydrofuran (20 mL) and water (20 mL). Purification by chromatography (SiO₂, 98:2 *isohexane*:ethyl acetate as
15 eluant) gave the sub-title compound as a solid (2.15 g).

¹H NMR (300 MHz, d₆-DMSO) δ 7.82 (1H, dd), 7.67 (1H, td), 7.61-7.52 (3H, m), 7.49-7.43 (2H, m), 3.64 (3H, s), 1.55 (9H, s).

20 **d) 4'-Chloro-[1,1'-biphenyl]-2,3'-dicarboxylic acid, 2-methyl ester**

Trifluoroacetic acid (3.3 mL) was added to a stirred solution of 4'-chloro-[1,1'-biphenyl]-2,3'-dicarboxylic acid, 3'-(1,1-dimethylethyl) 2-methyl ester (Example 49 (c)) (2.15 g) in dichloromethane (10 mL) and the mixture was stirred at room temperature under nitrogen for 90 minutes. The mixture was then evaporated to afford the sub-title compound as a
25 solid (1.7 g).

¹H NMR (400 MHz, d₆-DMSO) δ 7.82 (1H, dd), 7.69-7.64 (2H, m), 7.59 (1H, d), 7.55 (1H, td), 7.49-7.44 (2H, m), 3.63 (3H, s).

30 **e) 4'-Chloro-3'-[[2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl]amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid, methyl ester**

N,N-Dimethylformamide (1 drop) and oxalyl chloride (0.16 mL) were added to a stirred solution of 4'-chloro-[1,1'-biphenyl]-2,3'-dicarboxylic acid, 2-methyl ester (Example 49 (d)) (170 mg) in dichloromethane (2 mL) at 0 °C. The reaction was allowed to warm to room temperature, stirred under nitrogen for 2 hours, then evaporated to dryness. The residue was dissolved in dichloromethane (2 mL) and cooled to 0 °C. [2-(1-adamantyl)ethyl]amine hydrochloride (153 mg) was added followed by triethylamine (0.24 mL). The reaction was allowed to warm to room temperature and stirred under nitrogen for 2 hours then poured into saturated aqueous sodium bicarbonate (20 mL). The aqueous was extracted with dichloromethane (3 x 20 mL). The combined organic fractions were dried (MgSO₄), filtered and evaporated. Purification by chromatography (SiO₂, 1:4 ethyl acetate:isohexane as eluant) afforded the sub-title compound as a solid (260 mg).

MS: APCI(+ve) 452/454 (M+H⁺).

¹H NMR (400 MHz, d₆-DMSO) δ 8.36 (1H, t), 7.79 (1H, dd), 7.66 (1H, td), 7.56-7.49 (2H, m), 7.46 (1H, dd), 7.32 (1H, dd), 7.29 (1H, d), 3.63 (3H, s), 3.27-3.21 (2H, m), 1.93 (3H, s), 1.73-1.56 (6H, m), 1.51 (6H, s), 1.35-1.28 (2H, m).

f) 4'-Chloro-3'-[[[(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid

Prepared according to the method of Example 48 (c), using 4'-chloro-3'-[[[(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid, methyl ester (Example 49 (e)) (260 mg), potassium hydroxide (100 mg), water (1 mL), methanol (1 mL) and tetrahydrofuran (1 mL), to afford the title compound as a solid (140 mg).


MS: APCI(-ve) 436 (M-H⁺).

m.p. 114-117°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.36 (1H, t), 7.77 (1H, dd), 7.59 (1H, td), 7.52-7.46 (2H, m), 7.42-7.34 (2H, m), 7.32 (1H, d), 3.23 (2H, dt), 1.93 (3H, s), 1.67 (3H, d), 1.61 (3H, d), 1.51 (6H, s), 1.34-1.28 (2H, m).

Example 50

3-[4-Chloro-3-[[[2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl]amino]carbonyl]phenyl]-2-pyridinecarboxylic acid



a) 3-Iodo-2-pyridinecarboxylic acid, methyl ester

- 5 Butyllithium (32 mL, 2.5 M in hexanes) was added dropwise over 10 minutes to a solution of 2,2,6,6-tetramethylpiperidine (10.2 mL) in tetrahydrofuran (100 mL) at -78°C under nitrogen. The mixture was stirred at -78°C for 15 minutes and then picolinic acid (2.4 g) was added portionwise over 10 minutes. After a further 10 minutes at -78°C the mixture was allowed to warm to 0°C and stirred under nitrogen for 30 minutes. The reaction
- 10 mixture was then added dropwise over 15 minutes to a solution of iodine (15 g) in tetrahydrofuran (100 mL) at 0°C . This was then allowed to warm to room temperature and stirred for 1 hour before water (20 mL) was added. The mixture was evaporated to dryness to leave a black oil. Dichloromethane (50 mL) was added and the mixture was cooled to 0°C . *N,N*-Dimethylformamide (1 drop) and oxalyl chloride (4 mL) were added. The reaction
- 15 was allowed to warm to room temperature and stirred under nitrogen for 2 hours, then evaporated to dryness. The residue was dissolved in dichloromethane (20 mL) and then methanol (20 mL) was added. The mixture was then stirred for 10 minutes before being evaporated to afford the sub-title compound as an oil (1.0 g) which was used in the next step without purification.

20

MS: APCI(+ve) 264 ($\text{M}+\text{H}^+$).

b) 3-[4-Chloro-3-[(1,1-dimethylethoxy)carbonyl]phenyl]-2-pyridinecarboxylic acid, methyl ester

2-Chloro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-benzoic acid, 1,1-dimethylethyl ester (Example 49 (b)) (500 mg), 3-iodo-2-pyridinecarboxylic acid, methyl ester (Example 50 (a)) (400 mg) and tetrahydrofuran (2 mL) were placed in a 10 mL microwave vial. A solution of potassium carbonate (400 mg) in water (1 mL) was added followed by *bis*(triphenylphosphine)palladium(II) chloride (50 mg), and the mixture was heated to 130 °C in a microwave for 3 hours then concentrated. The residue was partitioned between dichloromethane (20 mL) and water (20 mL). The layers were separated and the aqueous was extracted with dichloromethane (2x20 mL). The combined organics were dried, filtered and evaporated. Purification by chromatography (SiO₂, 1:4 ethyl acetate:isohexane as eluant) gave the sub-title compound as a solid (240 mg).

MS: APCI(+ve) 348/450 (M+H⁺).

¹H NMR (400 MHz, d₆-DMSO) δ 8.68 (1H, dd), 8.02 (1H, dd), 7.71-7.63 (3H, m), 7.54 (1H, dd), 3.71 (3H, s), 1.56 (9H, s).

c) 3-(3-Carboxy-4-chlorophenyl)-2-pyridinecarboxylic acid, 2-methyl ester

Prepared according to the method of Example 49 (d), using 3-[4-chloro-3-[(1,1-dimethylethoxy)carbonyl]phenyl]-2-pyridinecarboxylic acid, methyl ester (Example 50 (b)) (240 mg), trifluoroacetic acid (1 mL) and dichloromethane (3 mL) to afford the sub-title compound as an oil (200 mg).

MS: APCI(+ve) 292/294 (M+H⁺).

¹H NMR (300 MHz, CDCl₃) δ 8.97 (1H, dd), 8.14 (1H, dd), 8.00 (1H, d), 7.91 (1H, dd), 7.64 (1H, d), 7.48 (1H, dd), 3.88 (3H, d).

d) 3-[4-Chloro-3-[(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid, methyl ester

Prepared according to the method of Example 49 (e), using 3-(3-carboxy-4-chlorophenyl)-2-pyridinecarboxylic acid, 2-methyl ester (Example 50 (c)) (170 mg), *N,N*-dimethylformamide (1 drop), oxalyl chloride (0.16 mL), [2-(1-adamantyl)ethyl]amine

hydrochloride (150 mg) and dichloromethane (4 mL). Purification (SiO₂, 2 % methanol in dichloromethane) afforded the sub-title compound as a solid (200 mg).

MS: APCI(+ve) 453/455 (M+H⁺).

5 ¹H NMR (400 MHz, CDCl₃) δ 8.71 (1H, dd), 7.74 (1H, dd), 7.66 (1H, d), 7.51 (1H, dd), 7.45 (1H, d), 7.31 (1H, dd), 6.17 (1H, s), 3.85 (3H, s), 3.53-3.45 (2H, m), 1.97 (3H, s), 1.72 (3H, d), 1.64 (3H, d), 1.56 (6H, s), 1.45-1.38 (2H, m),

10 **e) 3-[4-Chloro-3-[[2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid**

Prepared according to the method of Example 48 (c), using 3-[4-chloro-3-[[2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid, methyl ester (Example 50 (d)) (200 mg), potassium hydroxide (100 mg), water (1 mL), methanol (1 mL) and tetrahydrofuran (1 mL). Purification (Varian NH₂ cartridge using
15 dichloromethane (100 mL) and then 5 % acetic acid in dichloromethane (100 mL) as eluant) afforded the title compound as a solid (85 mg).


MS: APCI(+ve) 439/441 (M+H⁺).

m.p. 108-111°C.

20 ¹H NMR (300 MHz, d₆-DMSO) δ 8.62 (1H, dd), 8.39 (1H, t), 7.92 (1H, dd), 7.65-7.52 (2H, m), 7.50-7.40 (2H, m), 3.30-3.14 (2H, m), 1.92 (3H, s), 1.75-1.45 (12H, m), 1.36-1.27 (2H, m).

Example 51

25 **4'-Chloro-4-fluoro-3'-[[2-tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid**



a) 2-Bromo-5-fluoro-benzoic acid, methyl ester

Prepared according to the method of Example 13 (a) using 2-bromo-5-fluoro-benzoic acid (160 mg), oxalyl chloride (0.2 mL), *N,N*-dimethylformamide (1 drop), dichloromethane (2 mL) and methanol (2 mL) to give the sub-title compound as a solid (170 mg).

^1H NMR (400 MHz, CDCl_3) δ 7.63 (1H, dd), 7.53 (1H, dd), 7.07 (1H, ddd), 3.94 (3H, s).

b) 4'-Chloro-4-fluoro-3'--[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid, methyl ester

[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (200 mg), 2-bromo-5-fluoro-benzoic acid, methyl ester (Example 51 (a)) (170 mg), tetrahydrofuran (2 mL), a solution of potassium carbonate (166 mg) in water (1 mL) and *bis*(triphenylphosphine)palladium(II) chloride (20 mg) were placed in a 10 mL microwave vial. The mixture was heated to 70 °C for 1 hour in a microwave then evaporated. The residue was partitioned between dichloromethane (20 mL) and water (20 mL). The layers were separated and the aqueous was extracted with dichloromethane (2x20 mL). The combined organics were dried (MgSO_4), filtered and evaporated. Purification by chromatography (SiO_2 , 1:4 ethyl acetate:*isohexane* as eluant) gave the sub-title compound as an oil (79 mg).

MS: APCI(+ve) 456/458 ($\text{M}+\text{H}^+$).

c) 4'-Chloro-4-fluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid

Prepared according to the method of Example 48 (c), using 4'-chloro-4-fluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid, methyl ester (Example 51 (b)) (79 mg), potassium hydroxide (50 mg), water (0.5 mL), methanol (0.5 mL) and tetrahydrofuran (0.5 mL) to afford the title compound as a solid (65 mg).


MS: APCI(-ve) 440 (M-H⁺).

m.p. 126-128°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.37 (1H, t), 7.56 (1H, d), 7.50 (1H, d), 7.48-7.43 (2H, m), 7.39-7.32 (2H, m), 2.94 (2H, d), 1.94 (3H, s), 1.67 (3H, d), 1.59 (3H, d), 1.52 (6H, s).

Example 52

2-[5-Chloro-4-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-2-pyridinyl]-benzoic acid



(a) 2-[5-Chloro-4-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-2-pyridinyl]-benzoic acid, ethyl ester

To a mixture of 2,5-dichloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-4-pyridinecarboxamide (Prepared as described in WO 01/94338) (250 mg), 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-benzoic acid, ethyl ester (204 mg) and potassium carbonate (100 mg) in tetrahydrofuran (3 mL) and water (0.5 mL) was added *tetrakis*(triphenylphosphine)palladium(0) (20 mg). The mixture was heated at 80°C under a

nitrogen atmosphere for 17 hours and subsequently cooled to room temperature and water (5 mL) added. The resulting solid was collected by filtration and washed with water (10 mL). Drying of the solid yielded the sub-title compound as a solid (250 mg).

5 MS: APCI(-ve) 451/453 ($M-H^+$).

(b) 2-[5-Chloro-4-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]-2-pyridinyl]-benzoic acid

To a solution of 2-[5-chloro-4-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]-2-pyridinyl]-benzoic acid, ethyl ester (Example 52 (a)) (250 mg) in methanol (4 mL) was
10 added aqueous sodium hydroxide solution (40%, 2 mL). The mixture was stirred at room temperature for 18 hours, acidified with 2M aqueous hydrochloric acid and the resulting solid removed by filtration. Recrystallisation (acetonitrile) yielded the title compound as a colourless solid (180 mg).


15 MS: APCI(+ve) 425/427 ($M+H^+$).

m.p. 134-138°C.

¹H NMR (400 MHz, d₆-DMSO) δ 12.82 (1H, s), 8.69 (1H, d), 8.65 (1H, t), 7.74 (1H, d), 7.66-7.61 (3H, m), 7.57 (1H, m), 2.97 (2H, d), 1.95 (3H, s), 1.72-1.56 (6H, m), 1.51-1.53
20 (6H, m).

Example 53

2-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-4-methyl-3-pyridinecarboxylic acid



a) 4-Methyl-3-pyridinecarboxylic acid, butyl ester

4-Methyl-3-pyridinecarboxylic acid hydrochloride (1.5 g) and n-butanol (6 mL) were heated in a microwave at 180 °C for 90 minutes. Saturated aqueous sodium hydrogen carbonate was added and the mixture was extracted three times with dichloromethane. The combined organics were washed with water, brine, dried over magnesium sulphate, filtered and the solvent removed to afford the sub-title compound (1.0 g).

MS: APCI(+ve) 194 (M+H⁺).

b) 4-Methyl-3-pyridinecarboxylic acid 1-oxide, butyl ester

To a stirred solution of 4-methyl-3-pyridinecarboxylic acid, butyl ester (Example 53 (a)) (1.0 g) in dichloromethane (5mL) was added 36-40% peracetic acid (1 mL). After stirring for 12 hours, saturated aqueous sodium hydrogen carbonate was added and the mixture was extracted three times with dichloromethane. The combined organics were washed with water, brine, dried over magnesium sulphate, filtered and the solvent removed to afford the sub-title compound as an oil (1.0 g).

MS: APCI(+ve) 210 (M+H⁺).

c) 2-Chloro-4-methyl-3-pyridinecarboxylic acid, butyl ester

To 4-methyl-3-pyridinecarboxylic acid 1-oxide, butyl ester (Example 53 (b)) (1.0 g) was added phosphorus oxychloride (2 mL) which was then heated at 80 °C for 5 hours. The volatile components were removed under reduced pressure, ice was then added and the

mixture was stirred for 2 hours. The mixture was extracted three times with dichloromethane and the combined organics were washed with saturated aqueous sodium hydrogen carbonate, brine, dried over magnesium sulphate, filtered and the solvent removed to afford the sub-title compound as a brown oil (500 mg).

MS: APCI(+ve) 228 ($M+H^+$).

d) 2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-methyl-3-pyridinecarboxylic acid

2-Chloro-4-methyl-3-pyridinecarboxylic acid, butyl ester (Example 53 (c)) (250 mg), [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (347 mg), sodium carbonate (318 mg), *tetrakis*(triphenylphosphine)palladium(0) (20 mg), tetrahydrofuran (2 mL), and water (1 mL) were heated in a microwave for 90 minutes at 120 °C. 48% sodium hydroxide solution (200 µl) was added and the mixture was heated in a microwave at 90 °C for 60 minutes. The products were acidified with 2M aqueous hydrochloric acid, extracted three times with ethyl acetate, the combined organics were washed with brine, dried over magnesium sulphate, filtered and the solvent removed *in vacuo*. Purification by chromatography (SiO₂, dichloromethane:methanol:acetic acid 96.5:3:0.5 as eluant) and then by trituration with diethyl ether afforded the title compound as a solid (19 mg).

MS: APCI(+ve) 439 ($M+H^+$).


m.p. 177-180 °C.

¹H NMR (400 MHz, d₆-DMSO) δ 13.45 (1H, s), 8.62 (1H, d), 8.27 (1H, t), 7.41 - 7.38 (1H, m), 7.33 (1H, d), 7.26 - 7.22 (2H, m), 4.39 (3H, s), 2.90 (2H, d), 1.93 (3H, s), 1.70 - 1.47 (12H, m).

Example 54

6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-[(2-hydroxyethyl)methylamino]-3-pyridinecarboxylic acid

129



a) 6-Chloro-2-[(2-hydroxyethyl)methylamino]-3-pyridinecarboxylic acid

2,6-Dichloro-3-pyridinecarboxylic acid (500 mg) and 2-(methylamino)-ethanol (586 mg) were stirred in acetonitrile (2 mL) for 12 hours and then heated in a microwave at 70 °C for 60 minutes. The solvent was removed under vacuum to afford the sub-title compound (600 mg).

MS: APCI(-ve) 229 (M-H⁺).

b) 6-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-2-[(2-hydroxyethyl)methylamino]-3-pyridinecarboxylic acid

6-Chloro-2-[(2-hydroxyethyl)methylamino]-3-pyridinecarboxylic acid (Example 54 (a)) (240 mg), [4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-boronic acid (Example 2(a)) (200 mg), sodium carbonate (122 mg),

tetrakis(triphenylphosphine)palladium (0) (20 mg), tetrahydrofuran (1 mL), and water (1 mL) were heated in a microwave for 30 minutes at 120 °C. The reaction was acidified with 2M aqueous hydrochloric acid and extracted three times with ethyl acetate. The combined organics were washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. Purification by RP-HPLC (acetonitrile: aqueous trifluoroacetic acid, Symmetry) and then by trituration with diethyl ether afforded the title compound as a solid (72 mg).


MS: APCI(+ve) 498 (M+H⁺).

m.p. 121-124°C.

^1H NMR (300 MHz, d_6 -DMSO) δ 8.44 (1H, t), 8.14 - 8.06 (2H, m), 7.96 (1H, d), 7.59 (1H, d), 7.40 - 7.32 (1H, m), 3.70 - 3.58 (4H, m), 3.02 (3H, s), 2.97 (2H, d), 1.95 (3H, s), 1.73 - 1.52 (12H, m).

5 **Example 55**

3-[4-Methyl-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid



a) 5-Iodo-2-methyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

10 5-Iodo-2-methyl-benzoic acid (3.0 g) was stirred in dichloromethane (40 mL) under nitrogen. Oxalyl chloride (5 mL) was added followed by *N,N*-dimethylformamide (1 drop). After 2 hours the volatiles were removed under vacuum and the residue was redissolved in dichloromethane (40 mL). Tricyclo[3.3.1.1^{3,7}]decane-1-methanamine (2.23 mL) and triethylamine (3.18 mL) were added and the mixture was stirred under nitrogen for 2 hours.

15 2M aqueous hydrochloric acid was added, the layers were separated and the aqueous fraction was extracted twice with dichloromethane. The combined organics were washed with water, brine, dried (MgSO_4), filtered and concentrated *in vacuo*. Purification by trituration with diethyl ether afforded the sub-title compound (4.7 g).

20 MS: APCI(+ve) 410 ($\text{M}+\text{H}^+$).

b) 2-Methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

5-Iodo-2-methyl-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 55 (a)) (500 mg), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi-1,3,2-dioxaborolane (496 mg), potassium acetate (539 mg), *tetrakis*(triphenylphosphine)palladium(0) (20 mg), and *N,N*-dimethylformamide (2 mL) were heated at 90 °C for 60 minutes in a microwave. Ethyl acetate and water were added, the layers were separated and the aqueous phase was extracted twice with ethyl acetate. The combined organics were washed with water, brine, dried (MgSO₄), filtered and concentrated *in vacuo*. Purification by chromatography (SiO₂, dichloromethane as eluant) afforded the sub-title compound (292 mg).

MS: APCI(+ve) 410 (M+H⁺).

c) 3-[4-Methyl-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]- 2-pyridinecarboxylic acid

2-Methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 55 (b)) (146 mg), 3-iodo-2-pyridinecarboxylic acid, methyl ester (Example 50 (a)) (113 mg), sodium carbonate (113 mg), *tetrakis*(triphenylphosphine)palladium(0) (10 mg), tetrahydrofuran (1 mL) and water (1 mL) were heated in a microwave at 120 °C for 60 minutes. 48% sodium hydroxide solution (300 µL) was added and the mixture was heated for 30 minutes at 100 °C in a microwave. The reaction mixture was acidified with 2M aqueous hydrochloric acid, the layers were separated and the aqueous fraction was extracted three times with ethyl acetate. The combined organic layers were washed with water and the solvent was removed under vacuum. Purification by chromatography (SiO₂, dichloromethane:methanol:acetic acid 95:4:1 as eluant) and then by trituration with diethyl ether afforded the title compound (63 mg).


MS: APCI(+ve) 405 (M+H⁺).

m.p. 181-183°C.

¹H NMR (400 MHz, d₆-DMSO) δ 13.35 (1H, s), 8.59 (1H, dd), 8.19 (1H, t), 7.93 (1H, dd), 7.61 (1H, dd), 7.40 - 7.37 (2H, m), 7.33 (1H, d), 2.95 (2H, d), 2.39 (3H, s), 1.94 (3H, s), 1.72-1.55 (6H, m), 1.52 - 1.49 (6H, m).

Example 56

4-[4-Chloro-3-[[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-1,3-dimethyl-1*H*-pyrazole-5-carboxylic acid



Prepared according to the method of Example 55 (c) using [4-chloro-3-
[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2(a))
(139 mg) and 4-bromo-1,3-dimethyl-1*H*-pyrazole-5-carboxylic acid, methyl ester (140
mg). Purification by chromatography (SiO₂, dichloromethane:methanol:acetic acid
98.5:1:0.5 as eluant) gave the title compound (25 mg).


MS: APCI(+ve) 442 (M+H⁺).

m.p. 207°C.

¹H NMR (300 MHz, d₆-DMSO) δ 13.30 (1H, s), 8.35 (1H, t), 7.48 (1H, d), 7.33 (1H, dd), 7.26 (1H, d), 4.02 (3H, s), 2.93 (2H, d), 2.09 (3H, s), 1.97 - 1.90 (3H, m), 1.71 - 1.49 (12H, m).

Example 57

2-[4-Methyl-3-[[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid



2-Methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 55 (b)) (90 mg), 2-chloro-3-pyridinecarboxylic acid, ethyl ester (53 mg), sodium carbonate (70 mg), *tetrakis*(triphenylphosphine)palladium(0) (15 mg), tetrahydrofuran (1 mL) and water (1 mL) were heated in a microwave at 120 °C for 100 minutes. 48% sodium hydroxide solution (300 µL) was added and the mixture was heated for 60 minutes at 110 °C in a microwave. The reaction was acidified with 2M aqueous hydrochloric acid, the layers were separated and the aqueous phase was extracted three times with ethyl acetate. The combined organic layers were washed with water and the solvent was removed under vacuum. Purification by RP-HPLC, (acetonitrile:aqueous trifluoroacetic acid, Symmetry) gave the title compound (60 mg).


MS: APCI(+ve) 405 (M+H⁺).

¹H NMR (300 MHz, d₆-DMSO) δ 8.76 (1H, dd), 8.20 (1H, t), 8.11 (1H, dd), 7.64 - 7.45 (3H, m), 7.31 (1H, d), 2.95 (2H, d), 2.40 (3H, s), 1.94 (3H, s), 1.63 (6H, q), 1.53 - 1.48 (6H, m).

Example 58

3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-2-pyridineacetic acid, monosodium salt

134



[4-Chloro-3-[[tricyclo[3.3.1.1.^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-boronic acid (Example 2(a)) (142 mg), 3-bromo-2-pyridineacetic acid ethyl ester (Prepared according to the method of Synthesis, 1997, 949-952) (100 mg), sodium carbonate (130 mg),
 5 *tetrakis*(triphenylphosphine)palladium(0) (10 mg), tetrahydrofuran (2 mL) and water (1 mL) were heated in a microwave at 120 °C for 30 minutes. 48% sodium hydroxide solution (200 µL) was added to the reaction which was stirred for 12 hours before being filtered. The solid was washed with water (3 mL) and then acetonitrile (3 mL) before being dried in a vacuum oven to afford the title compound as a solid (55 mg).

10 MS: APCI(-ve) 437 ($M-H^+$).


m.p. 186-187°C dec.

¹H NMR (300 MHz, d₆-DMSO) δ 8.61 (1H, t), 8.44 (1H, d), 7.72 (1H, dd), 7.61 - 7.55 (2H, m), 7.49 (1H, d), 7.26 - 7.19 (1H, m), 3.29 (2H, s), 2.94 (2H, d), 1.93 (3H, s), 1.70 -
 15 1.50 (12H, m).

Example 59

1-[3-[4-Chloro-3-[[tricyclo[3.3.1.1.^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-2-pyridinyl]-4-piperidinecarboxylic acid

135



a) **1-(3-Bromo-2-pyridinyl)-4-piperidinecarboxylic acid, ethyl ester**

A mixture of 2,3-dibromopyridine (250 mg), ethyl isonipecotate (250 mg) and triethylamine (0.25 mL) in acetonitrile (0.5 mL) was heated at 130°C in a microwave for 3 hours. The mixture was then concentrated to dryness *in vacuo* and partitioned between dichloromethane and water. The organics were collected, dried over magnesium sulphate, filtered and concentrated to dryness to give the sub-title compound as a brown oil (325 mg).

MS: APCI(+ve) 313/315 ($M+H^+$).

b) **1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-4-piperidinecarboxylic acid, ethyl ester**

A mixture of [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (300 mg), 1-(3-bromo-2-pyridinyl)-4-piperidinecarboxylic acid, ethyl ester (Example 59 (a)) (320 mg), potassium carbonate (240 mg) and *bis*(triphenylphosphine)palladium(II) chloride (60 mg) in toluene (10 mL) / ethanol (1 mL) / water (1 mL) was heated at 50°C under a nitrogen atmosphere for 3 hours. The mixture was concentrated *in vacuo*, passed through a plug of silica, eluting with ethyl acetate, and purified further by chromatography (SiO₂, *isohexane*:ethyl acetate 7:3 as eluant) to give the sub-title compound as a foam (160 mg).

MS: APCI(+ve) 536/538 ($M+H^+$).

c) 1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-4-piperidinecarboxylic acid

To a solution of 1-[3-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-

ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-4-piperidinecarboxylic acid, ethyl ester (Example 59 (b)) (160 mg) in tetrahydrofuran (2 mL) and ethanol (1 mL) was added a solution of potassium hydroxide (220 mg) in water (1 mL). The resulting solution was stirred at room temperature for 16 hours. The reaction mixture was then acidified by the addition of acetic acid and purified (Varian NH₂ cartridge using acetonitrile and then 50% acetic acid in acetonitrile as eluant) to give the title compound (90 mg).


MS: APCI(+ve) 508/510 (M+H⁺).

m.p. 110-115°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.37 (1H, t), 8.19 (1H, dd), 7.72 (1H, dd), 7.60 (1H, d), 7.57 (1H, dd), 7.54 (1H, d), 6.99 (1H, dd), 3.37 (2H, d), 2.95 (2H, d), 2.64 (2H, t), 1.80-1.38 (20H, m).

Example 60

1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-L-proline



a) 1-(3-Bromo-2-pyridinyl)-L-proline, 1,1-dimethylethyl ester

Prepared according to the method of Example 59 (a) using 2,3-dibromopyridine (250 mg) and L-proline 1,1-dimethylethyl ester (250 mg) to give the sub-title compound as an oil (325 mg).

¹H NMR (400 MHz, CDCl₃) δ 8.03 (1H, dd), 7.66 (1H, dd), 6.51 (1H, dd), 4.65 (1H, dd), 3.87-4.01 (2H, m), 2.18-2.33 (1H, m), 1.90-2.13 (3H, m), 1.40 (9H, s).

b) 1-[3-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-L-proline, 1,1-dimethylethyl ester

Prepared according to the method of Example 59 (b) using [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (300 mg) and 1-(3-bromo-2-pyridinyl)-L-proline, 1,1-dimethylethyl ester (Example 60 (a)) (320 mg) to give the sub-title compound as a foam (100 mg)

MS: APCI(+ve) 550/552 (M+H⁺)

c) 1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-L-proline

To a solution of 1-[3-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-L-proline, 1,1-dimethylethyl ester (Example 60 (b)) (200 mg) in dichloromethane (5 mL) was treated with trifluoroacetic acid (1 mL) and heated at reflux for 1 hour. The mixture was neutralised with 7N methanolic ammonia, concentrated to dryness *in vacuo* and the residue purified (Varian NH₂ cartridge using acetonitrile and then 20% acetic acid in acetonitrile as eluant) to give the title compound as a pale yellow solid (110 mg).


MS: APCI(+ve) 494 (M+H⁺).

m.p. 121-125°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.52 (1H, s), 8.06 (1H, dd), 7.51 (1H, d), 7.40-7.47 (3H, m), 6.76 (1H, dd), 4.34-4.43 (1H, m), 2.88-3.03 (3H, m), 2.71-2.78 (1H, m), 2.10-2.19 (1H, m), 1.93 (3H, s), 1.67-1.81 (3H, m), 1.66 (3H, d), 1.59 (3H, d), 1.53 (6H, s).

Example 61

1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-3-piperidinecarboxylic acid



a) **1-(3-Bromo-2-pyridinyl)-3-piperidinecarboxylic acid, ethyl ester**

Prepared according to the method of Example 59 (a) using 2,3-dibromopyridine (250 mg) and ethyl nipecotate (250 mg) to give the sub-title compound as an oil (325 mg).

MS: APCI(+ve) 313/315 ($M+H^+$).

b) **1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-3-piperidinecarboxylic acid, ethyl ester**

Prepared according to the method of Example 59 (b) using [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (300 mg) and 1-(3-bromo-2-pyridinyl)-3-piperidinecarboxylic acid, ethyl ester (Example 61 (a)) (320 mg) to give the sub-title compound as a foam (70 mg).

MS: APCI(+ve) 536/538 ($M+H^+$).

c) **1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-3-piperidinecarboxylic acid**

Prepared according to the method of Example 59 (c) using 1-[3-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-4-

piperidinecarboxylic acid, ethyl ester (Example 61 (b)) (70 mg) to give the title compound as a solid (35 mg).


MS: APCI(+ve) 508/510 ($M+H^+$).

5 m.p. 125-130°C.

1H NMR (400 MHz, d_6 -DMSO) δ 8.37 (1H, t), 8.19 (1H, dd), 7.72 (1H, dd), 7.60 (1H, d), 7.57 (1H, dd), 7.54 (1H, d), 6.99 (1H, dd), 3.37 (2H, d), 2.95 (2H, d), 2.64 (2H, t), 1.94 (3H, s), 1.67 (3H, d), 1.60 (3H, d), 1.53 (6H, s), 1.38-1.80 (m, 5H).

10 **Example 62**

1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-3-azetidinecarboxylic acid



a) **1-(3-Bromo-2-pyridinyl)-3-azetidinol**

15 To a solution of 2,3-dibromopyridine (700 mg) in pyridine (1 mL) at reflux was added, portionwise over 8 hours, azetidinol hydrochloride (1.4 g) and the mixture heated at reflux for a further 16 hours. The mixture was poured into water and the solid removed by filtration. The aqueous was then concentrated to dryness in vacuo and the residue purified (Varian C-18 cartridge, water : methanol gradient as eluant) to give the sub-title compound
20 as a solid (145 mg).

MS: APCI(+ve) 229/231 ($M+H^+$).

b) **1-(3-Bromo-2-pyridinyl)-3-azetidinol, methanesulfonate ester**

To a solution of 1-(3-bromo-2-pyridinyl)-3-azetidinol (Example 62 (a)) (145 mg) and triethylamine (0.1 mL) in dichloromethane (10 mL) was added dropwise a solution of methanesulfonyl chloride (0.07 mL) and the reaction mixture stirred at room temperature for 2 hours. The mixture was washed with dilute acetic acid and aqueous sodium bicarbonate solution. The organics were dried over magnesium sulphate, filtered, concentrated to dryness *in vacuo* and purified (SiO₂, dichloromethane as eluant) to give the sub-title compound as a colourless oil (194 mg).

MS: APCI(+ve) 307/309 (M+H⁺).

c) 1-(3-Bromo-2-pyridinyl)-3-azetidinedicarbonitrile

A mixture of 1-(3-bromo-2-pyridinyl)-3-azetidinol, methanesulfonate ester (Example 62 (b)) (194 mg) and sodium cyanide (388 mg) in N,N-dimethylformamide (2 mL) was heated at 110°C for 6 days. The reaction mixture was partitioned between water and dichloromethane. The organics were collected and purified by chromatography (SiO₂, dichloromethane as eluant) to give the sub-title compound as a colourless oil (115 mg).

MS: APCI(+ve) 238/240 (M+H⁺).

d) 1-(3-Bromo-2-pyridinyl)-3-azetidinedicarboxylic acid

A mixture of 1-(3-bromo-2-pyridinyl)-3-azetidinedicarbonitrile (Example 62 (c)) (150 mg) and potassium hydroxide (150 mg) in ethanol (2 mL) and water (2 mL) was heated to 100°C by microwave irradiation. After 30 minutes, the reaction was cooled and concentrated to dryness *in vacuo*. The residue was dissolved in water (5 mL) and extracted with dichloromethane (5 x 5 mL), the combined organic fractions were dried over magnesium sulphate, filtered and concentrated to dryness *in vacuo* to give the sub-title compound as an oil (150 mg).

MS: APCI(+ve) 257/259 (M+H⁺).

e) 1-(3-Bromo-2-pyridinyl)-3-azetidinedicarboxylic acid, methyl ester

To a solution of 1-(3-bromo-2-pyridinyl)-3-azetidinecarboxylic acid (Example 62 (d)) (150 mg) in dichloromethane (2 mL) containing *N,N*-dimethylformamide (1 drop), was added oxalyl chloride (0.5 mL) and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was concentrated to dryness and the residue dissolved in methanol (5 mL). The solution was stirred at room temperature for 30 minutes and then concentrated to dryness and the residue purified (Varian C-18 cartridge, water : methanol gradient as eluant) to give the sub-title compound as an oil (76 mg).

MS: APCI(+ve) 271/273 ($M+H^+$).

f) 1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-3-azetidinecarboxylic acid, methyl ester

Prepared according to the method of Example 59 (b) using [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (100 mg) and 1-(3-bromo-2-pyridinyl)-3-azetidinecarboxylic acid, methyl ester (Example 62 (e)) (74 mg) in tetrahydrofuran (2 mL) and water (2 mL) to give the sub-title compound as a foam (107 mg).

MS: APCI(+ve) 494/496 ($M+H^+$).

g) 1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-3-azetidinecarboxylic acid

Prepared according to the method of Example 59 (c) using 1-[3-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-3-azetidinecarboxylic acid, methyl ester (Example 62 (f)) (105 mg) to give the title compound as a solid (76 mg).


MS: APCI(+ve) 480 ($M+H^+$).

m.p. 145-150°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.36 (1H, t), 8.17 (1H, s), 7.33-7.56 (4H, m), 6.87 (1H, t), 3.78 (2H, t), 3.68 (2H, t), 3.26-3.39 (1H, m), 2.96 (2H, d), 1.94 (3H, s), 1.67 (3H, d), 1.60 (3H, d), 1.53 (6H, s).

Example 63

3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-6-methyl-2-pyridinecarboxylic acid



a) 2-Chloro-5-(2-chloro-6-methyl-3-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

A mixture of [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (174 mg), 3-bromo-2-chloro-6-methyl-pyridine (104 mg), potassium carbonate (138 mg) and *bis*(triphenylphosphine)palladium(II) chloride (27 mg) in tetrahydrofuran (2 mL) and water (2 mL) was stirred at room temperature under a nitrogen atmosphere over 72 hours. The mixture was filtered through diatomaceous earth, washing with methanol, and the filtrate was concentrated *in vacuo*. Purification by chromatography (SiO₂, *isohexane*:ethyl acetate 1:1 as eluant) gave the sub-title compound as a solid (135 mg).

MS: APCI(+ve) 429/431 (M+H⁺)

b) 2-Chloro-5-(2-cyano-6-methyl-3-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

A mixture of 2-chloro-5-(2-chloro-6-methyl-3-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 63 (a)) (0.13 g), *bis*(dibenzylideneacetone)palladium (0.05 g), 1,1'-*bis*(diphenylphosphino)ferrocene (0.11 g) and copper (I) cyanide (0.13 g) in 1,4-

dioxane (3 mL) was heated at reflux under nitrogen for 3 hours. The reaction was passed through a plug of silica, eluting with acetonitrile and the liquors concentrated to dryness *in vacuo*. The residue was then purified by chromatography (SiO₂, 1:2 ethyl acetate: isohexane as eluant) to give the sub-title compound as a solid (0.10 g)

MS: APCI(+ve) 420/422 (M+H⁺).

c) 3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-6-methyl-2-pyridinecarboxylic acid

A solution of 2-chloro-5-(2-cyano-6-methyl-3-pyridinyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 63 (b)) (70 mg) in acetonitrile (2 mL) was treated with a solution of potassium hydroxide (0.25 g) in water (0.6 mL) and the mixture was heated in a microwave at 120°C for 60 minutes. The reaction mixture was then concentrated to dryness *in vacuo*, dissolved in water (5 mL) and concentrated hydrochloric acid (5 mL) then heated at reflux for 24 hours. The reaction mixture was concentrated to dryness and purified (Varian C-18 cartridge, water:methanol gradient as eluant, then by Varian NH₂ cartridge using methanol (100 mL) and then 5 % acetic acid in methanol (100 mL) as eluant) to give the title compound as a solid (25 mg).

MS: APCI(+ve) 439 (M+H⁺).


m.p. 125-130°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.38 (1H, t), 7.75 (1H, d), 7.73 - 7.66 (1H, m), 7.50-7.47 (1H, m), 7.44 (1H, s), 7.39 (1H, d), 2.94 (2H, d), 2.07 (3H, s), 1.93 (3H, s), 1.70-1.56 (6H, m), 1.52 (6H, s).

Example 64

3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-5-methyl-2-pyridinecarboxylic acid

144



a) 3-Bromo-5-methyl-2-pyridinamine

To a solution of 2-amino-3-picoline (10.8 g) in glacial acetic acid (30 mL) at 80°C was added dropwise a solution of bromine (5.5 mL) in glacial acetic acid (5.5 mL) and the temperature maintained for 1 hour. The reaction was concentrated to dryness and the residue dissolved in water (100 mL) which was then basified with 0.880 ammonia solution and extracted into dichloromethane. The organics were dried (MgSO₄), filtered and concentrated to dryness. The resulting residue was recrystallised from *isohexane*/ethyl acetate to give the sub-title compound as a solid (18.3 g).

MS: APCI(+ve) 187/189 (M+H⁺).

b) 3-Bromo-2-chloro-5-methyl-pyridine

To a solution of 3-bromo-5-methyl-2-pyridinamine (Example 64 (a)) (1.0 g) in a mixture of concentrated hydrochloric acid (5 mL) and water (3 mL) at 0°C was added a solution of sodium nitrite (0.36 g) in water (3 mL). After the addition was complete, the reaction mixture was neutralised by the addition of 0.880 ammonia and the resulting precipitate collected by filtration and purified by chromatography (SiO₂, dichloromethane as eluant) to give the sub-title compound as a solid (0.47 g)

MS: APCI(+ve) 206 (M+H⁺).

c) 2-Chloro-5-(2-chloro-5-methyl-3-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

Prepared according to the method of Example 63 (a) using [4-chloro-3-
[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a))
(113 mg), 3-bromo-2-chloro-5-methyl-pyridine (Example 64 (b)) (209 mg) and
tetrakis(triphenylphosphine)palladium(0) (35 mg) at 50°C to give the sub-title compound
5 as a solid (220 mg).

MS: APCI(+ve) 429/431 (M+H⁺).

**d) 2-Chloro-5-(2-cyano-5-methyl-3-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
10 benzamide**

Prepared according to the method of Example 63 (b) using 2-chloro-5-(2-chloro-5-methyl-
3-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 64 (c)) (0.15 g) to
give the sub-title compound as a solid (0.13 g).

15 MS: APCI(+ve) 420/422 (M+H⁺).

**e) 3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-5-
methyl-2-pyridinecarboxylic acid**

Prepared according to the method of Example 21(c) using 2-chloro-5-(2-cyano-5-methyl-
20 3-pyridinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide (Example 64 (d)) (0.12 g)
to give the title compound as a solid (80 mg).

MS: APCI(+ve) 439 (M+H⁺).


m.p. 125-130°C.

25 ¹H NMR (300 MHz, CDCl₃) δ 8.44 (1H, s), 7.63 (1H, s), 7.59 (1H, s), 7.44 (1H, d), 7.35
(1H, d), 6.38 (1H, s), 3.16 (2H, d), 2.47 (3H, s), 2.00 (3H, s), 1.73 (3H, d), 1.64 (3H, d),
1.59 (6H, s).

Example 65

30 **1-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-
pyridinyl]-4-hydroxy-4-piperidinecarboxylic acid**

146



a) 1-(3-Bromo-2-pyridinyl)-4-hydroxy-4-piperidinecarboxylic acid, methyl ester

Prepared according to the method of Example 59 (a) using 2,3-dibromopyridine (145 mg) and 4-hydroxy-4-piperidinecarboxylic acid methyl ester (97 mg) to give the sub-title compound as an oil (44 mg).

MS: APCI(+ve) 315/317 ($M+H^+$).

b) 1-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-2-pyridinyl]-4-hydroxy-4-piperidinecarboxylic acid

Prepared according to the method of Example 59 (b) using [4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (55 mg) and 1-(3-bromo-2-pyridinyl)-4-hydroxy-4-piperidinecarboxylic acid, methyl ester (Example 65 (a)) (44 mg) in tetrahydrofuran (4 mL) and water (2 mL). Upon completion of the reaction potassium hydroxide (100 mg) in water (1 mL) was added and the mixture stirred at room temperature overnight. The reaction mixture was concentrated to dryness and the residue purified (Varian C-18 cartridge, water : methanol gradient as eluant) to give the title compound as a white solid (50 mg).


MS: APCI(+ve) 522 ($M+H^+$).

m.p. 145-150°C.

¹H NMR (300 MHz, CDCl₃) δ 8.27 (1H, d), 8.21 (1H, d), 7.60 (1H, dd), 7.51 (1H, dd), 7.49 (1H, d), 6.96 (1H, dd), 6.49 (1H, t), 3.37 (2H, d), 3.22 - 3.11 (2H, m), 3.20 (2H, d), 2.14 - 1.96 (4H, m), 2.01 (3H, s), 1.74 (3H, d), 1.64 (3H, d), 1.59 (6H, s).

Example 66

1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-5-fluoro-2-pyridinyl]-4-piperidinecarboxylic acid



a) 3-Bromo-5-fluoro-2-pyridinamine

Prepared according to the method of Example 64 (a) using 5-fluoro-2-pyridinamine (2.0 g) to give the sub-title compound as a solid (2.2 g).

MS: APCI(+ve) 191/193 ($M+H^+$).

b) 2,3-Dibromo-5-fluoro-pyridine

To a solution of 3-bromo-5-fluoro-2-pyridinamine (Example 66 (a)) (1.0 g) in hydrobromic acid (2.5 mL) at 0°C was added bromine (0.85 mL) dropwise, maintaining the temperature below 5°C. Then a solution of sodium nitrite (0.92 g) in water (2 mL) was added dropwise again at below 5°C. The mixture was then stirred at 0°C for 30 minutes before being treated dropwise with a solution of sodium hydroxide (2.0 g) in water (2 mL). The reaction mixture was allowed to warm to room temperature and was partitioned between water and ethyl acetate, the organics were washed with water, dried ($MgSO_4$), filtered and concentrated to dryness to give an orange oil which was purified by chromatography (SiO_2 cartridge eluting with dichloromethane) to give the sub-title compound as a solid (1.1 g).

1H NMR (300 MHz, $CDCl_3$) δ 7.71 (1H, dd), 8.26 (1H, d).

c) **1-(3-Bromo-5-fluoro-2-pyridinyl)-4-piperidinecarboxylic acid, methyl ester**

Prepared according to the method of Example 59 (a) using 2,3-dibromo-5-fluoro-pyridine (Example 66 (b)) (250 mg) and methyl isonipecotate (250 mg) to give the sub-title compound as an oil (125mg).

5

MS: APCI(+ve) 317/319 ($M+H^+$).

d) **1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-5-fluoro-2-pyridinyl]-4-piperidinecarboxylic acid, methyl ester**

10 Prepared according to the method of Example 26(a) using [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (146 mg) and 1-(3-bromo-5-fluoro-2-pyridinyl)-4-piperidinecarboxylic acid, methyl ester (Example 66 (c)) (120 mg) to give the sub-title compound as a solid (150 mg).

15 MS: APCI(+ve) 540/542 ($M+H^+$).

e) **1-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-5-fluoro-2-pyridinyl]-4-piperidinecarboxylic acid**

To a suspension of 1-[3-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-5-fluoro-2-pyridinyl]-4-piperidinecarboxylic acid, methyl ester (Example 66 (d)) (150 mg) in methanol (8 mL) was added a solution of sodium hydroxide (200 mg) in water (1.5 mL). The resulting mixture was then stirred at room temperature over 72 hours. The reaction was acidified with acetic acid, concentrated to dryness and purified (Varian NH₂ cartridge using acetonitrile and then 50% acetic acid
25 in acetonitrile as eluant) to give the title compound as a solid (114 mg).


MS: APCI(+ve) 526 ($M+H^+$).

m.p. 130-135°C.

1H NMR (300 MHz, d₆-DMSO) δ 8.34 (1H, t), 8.22 (1H, d), 7.77 (1H, d), 7.73 (1H, s),
30 7.64 (1H, dd), 7.58 (1H, d), 3.28 (2H, d), 2.95 (2H, d), 2.65 (2H, t), 2.27 (1H, t), 1.95 (3H, s), 1.68 (3H, d), 1.59 (3H, d), 1.53 (6H, s), 1.47-1.77 (4H, m).

Example 67

4'-Methyl-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid



2-Methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 55 (b)) (150 mg), 2-bromo-benzoic acid methyl ester (79 mg), sodium carbonate (116 mg), *tetrakis*(triphenylphosphine)palladium(0) (15 mg), tetrahydrofuran (2 mL) and water (1 mL) were heated in a microwave at 120 °C for 30 minutes. 48% Sodium hydroxide solution (300 µL) was added and the mixture was heated for 30 minutes at 120 °C in a microwave. The reaction was neutralised with acetic acid and purification by RP-HPLC (acetonitrile:aqueous trifluoroacetic acid, Symmetry) gave the title compound (58 mg).


MS: APCI(+ve) 404 (M+H⁺).

m.p. 85-92°C.

¹H NMR (300 MHz, d₆-DMSO) δ 8.16 (1H, t), 7.74 - 7.70 (1H, m), 7.62 - 7.55 (1H, m), 7.49 - 7.41 (2H, m), 7.33 - 7.23 (3H, m), 2.95 (2H, d), 2.38 (3H, s), 1.94 (3H, s), 1.71 - 1.55 (6H, m), 1.51 (6H, s).

Example 68

1-[3-[4-Methyl-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-4-piperidinecarboxylic acid



a) 1-(3-Bromo-2-pyridinyl)-4-piperidinecarboxylic acid methyl ester

2,3-Dibromopyridine (3.0 g) and methyl isonipecotate (5.4 g) were heated in a microwave at 130 °C for 30 minutes. Purification by chromatography (SiO₂, dichloromethane as eluant) gave the sub-title compound as a colourless oil (2.4 g).

MS: APCI(+ve) 299/301 (M+H⁺).

¹H NMR (400 MHz, d₆-DMSO) δ 8.24 (1H, dd), 7.95 (1H, dd), 6.92 (1H, dd), 3.63 (3H, s), 3.62-3.59 (2H, m), 2.84 (2H, dd), 2.60-2.52 (1H, m), 1.97-1.89 (2H, m), 1.77-1.66 (2H, m).

b) 1-[3-[4-Methyl-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-4-piperidinecarboxylic acid

2-Methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 55 (b)) (200 mg), 1-(3-bromo-2-pyridinyl)-4-piperidinecarboxylic acid methyl ester (Example 68 (a)) (146 mg), sodium carbonate (155 mg), *tetrakis*(triphenylphosphine)palladium(0) (15 mg), tetrahydrofuran (2 mL) and water (1 mL) were heated in a microwave at 120 °C for 60 minutes. 48% Sodium hydroxide solution (300 µL) and methanol (2mL) were added and the mixture stirred for 12 hours. The reaction was neutralised with acetic acid and purified by RP-HPLC, (acetonitrile:aqueous trifluoroacetic acid, Symmetry) to give the title compound (77 mg).


MS: APCI(+ve) 488 (M+H⁺).

m.p. 60°C.

^1H NMR (300 MHz, d_6 -DMSO) δ 8.21 - 8.12 (2H, m), 7.73 (1H, d), 7.57 (2H, d), 7.33 (1H, d), 7.10 (1H, dd), 3.43 (2H, d), 2.96 (2H, d), 2.75 (2H, t), 2.38 (3H, s), 2.36 - 2.26 (1H, m), 1.95 (3H, s), 1.79 - 1.42 (16H, m).

5 **Example 69**

6-[4-Methyl-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid



2-Methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 55 (b)) (200 mg), 6-chloro-2-pyridinecarboxylic acid, methyl ester (108 mg), sodium carbonate (207 mg), *tetrakis*(triphenylphosphine)palladium (0) (15 mg), tetrahydrofuran (2 mL) and water (1 mL) were heated in a microwave at 120 °C for 60 minutes. 48% Sodium hydroxide solution (300 μL) and methanol (2 mL) were added and the mixture was heated for 30 minutes at 90 °C in a microwave. The reaction
15 was neutralised with acetic acid and purified by RP-HPLC, (acetonitrile:aqueous trifluoroacetic acid, Symmetry) to give the title compound (44 mg).


MS: APCI(+ve) 405 ($\text{M}+\text{H}^+$).

m.p. 227°C.

20 ^1H NMR (400 MHz, d_6 -DMSO) δ 13.20 (1H, s), 8.26 (1H, t), 8.21 (1H, dd), 8.14 - 8.10 (2H, m), 8.07 (1H, t), 7.98 (1H, dd), 7.39 (1H, d), 2.99 (2H, d), 2.41 (3H, s), 1.95 (3H, s), 1.71 - 1.59 (6H, m), 1.56 - 1.52 (6H, m).

Example 70

4-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid



A mixture of [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (200 mg), 4-chloronicotinic acid (91 mg), potassium carbonate (159 mg) and *tetrakis*(triphenylphosphine)palladium(0) (67 mg) in tetrahydrofuran (0.5 mL) and water (0.5 mL) were heated in a microwave at 120°C for 2 hours. The products were filtered through diatomaceous earth, washing with methanol (2 x 10 mL), the filtrate was concentrated and the residue purified by RP-HPLC (acetonitrile :aqueous trifluoroacetic acid, Symmetry) to give the title compound as a solid (7 mg).


MS: APCI(+ve) 425 (M+H⁺).

m.p. 104-107°C.

¹H NMR (300 MHz, d₆-DMSO) δ 8.95 (1H, s), 8.77 (1H, d), 8.41 (1H, t), 7.58 (1H, d), 7.51-7.42 (3H, m), 2.95 (2H, d), 1.94 (3H, s), 1.72 - 1.56 (6H, m), 1.53 (6H, s).

Example 71

6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-methyl-2-pyridinecarboxylic acid



a) 3-Methyl-2-pyridinecarboxylic acid 1-oxide, methyl ester

A solution of 3-methyl 2-pyridinecarboxylic acid, methyl ester (340 mg) in acetic acid (5 mL) and 35% aqueous hydrogen peroxide (5 mL) was heated at 60°C for 5 hours before
 5 being stirred at room temperature overnight. The reaction mixture was quenched by pouring into a sodium sulfite/ ice water mixture (50 mL) and then extracted into dichloromethane (3 x 25 mL). The organics were dried over magnesium sulphate, filtered and concentrated to dryness to give the sub-title compound as a colourless oil (300 mg).

10 MS: APCI(+ve) 168 ($M+H^+$).

b) 6-Chloro-3-methyl-2-pyridinecarboxylic acid, methyl ester

A solution of 3-methyl-2-pyridinecarboxylic acid 1-oxide, methyl ester (Example 71 (a)) (300 mg) in phosphorous oxychloride (1 mL) was heated at 60°C for 16 hours. The
 15 reaction mixture was then poured into water (10 mL) and extracted with dichloromethane (2 x 10 mL). The organics were dried over magnesium sulphate, filtered and concentrated to dryness to give the sub-title compound as a colourless oil (300 mg).

MS: APCI(+ve) 185/187 ($M+H^+$).

c) 6-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-3-methyl-2-pyridinecarboxylic acid

A mixture of [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (190 mg), 6-chloro-3-methyl-2-pyridinecarboxylic acid, methyl ester (Example 71(b)) (200 mg), potassium carbonate (150 mg) and dichlorobis(triphenylphosphine) palladium (II) (30 mg) in tetrahydrofuran (7 mL) and water (7 mL) were heated at reflux for 5 hours, sodium hydroxide (100 mg) was added and the mixture was stirred for 16 hours at room temperature. The products were concentrated and purified (Varian C-18 cartridge, water:acetonitrile gradient as eluant, followed by Varian NH₂ resin, acetonitrile:acetic acid gradient as eluant) to give the title compound as a solid (60 mg).


MS: APCI(+ve) 439 (M+H⁺).

m.p. 180-185°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.43 (1H, t), 8.19 - 8.15 (2H, m), 8.08 (1H, d), 7.86 (1H, d), 7.60 (1H, d), 2.98 (2H, d), 2.47 (3H, s), 1.95 (3H, s), 1.68 (3H, d), 1.61 (3H, d), 1.55 (6H, s).

Example 72

6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-(trifluoromethyl)-2-pyridinecarboxylic acid



a) 6-Chloro-4-(trifluoromethyl)-2-pyridinecarboxylic acid, methyl ester

To a solution of 6-chloro-4-trifluoromethoxy-pyridine-2-carboxylic acid (144 mg) in dichloromethane (5 mL) containing 1 drop of *N,N*-dimethylformamide was added dropwise

oxalyl chloride (0.2 mL). The mixture was stirred at room temperature for 1 hour and then concentrated to dryness *in vacuo*, azeotroping with dichloromethane (x 3). The residue was dissolved in methanol and the solution stirred at room temperature for 1 hour then concentrated to dryness in *vacuo* to give the sub-title compound as a white solid (152 mg).

5

¹H NMR (300 MHz, CDCl₃) δ 8.27 (1H, s), 7.76 (1H, s), 4.05 (3H, s).

b) 6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-(trifluoromethyl)-2-pyridinecarboxylic acid

10

Prepared according to the method of Example 71 (c) using [4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (230 mg), 6-chloro-4-(trifluoromethyl)-2-pyridinecarboxylic acid, methyl ester (Example 72 (a)) (152 mg), potassium carbonate (170 mg) and dichlorobis(triphenylphosphine) palladium (II) (20 mg) at 50°C to give the title compound as a solid (50 mg).

15

MS: APCI(+ve) 493 (M+H⁺).


m.p. 220-225°C.

¹H NMR (300 MHz, d₆-DMSO) δ 8.69 (1H, s), 8.46 (1H, t), 8.39 - 8.36 (2H, m), 8.21 (1H, s), 7.68 (1H, t), 2.99 (2H, d), 1.96 (3H, s), 1.69 (3H, d), 1.61 (3H, d), 1.57 (6H, s).

20

Example 73

5-[6-(Acetylamino)-2-methyl-3-pyridinyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



a) *N*-(5-Bromo-6-methyl-2-pyridinyl)-acetamide

To a solution of 5-bromo-6-methyl-2-pyridinamine (500 mg) in dichloromethane (10 mL) was added triethylamine (0.42 mL) followed by acetyl chloride (0.25 mL). The reaction was heated at reflux for 1 hour then cooled and washed with water. The organics were dried over magnesium sulphate, filtered and concentrated to dryness *in vacuo* and the residue was triturated with *isohexane* to give the sub-title compound as a solid (520 mg).

MS: APCI(+ve) 229/231 ($M+H^+$).

b) 5-[6-(Acetylamino)-2-methyl-3-pyridinyl]-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

A mixture of [4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-boronic acid (Example 2 (a)) (350 mg), *N*-(5-bromo-6-methyl-2-pyridinyl)-acetamide (Example 73 (a)) (229 mg), potassium carbonate (280 mg) and dichlorobis(triphenylphosphine)palladium (II) (50 mg) in toluene (10 mL) / water (1 mL) and ethanol (1 mL) was heated at 80°C under a nitrogen atmosphere for 3 hours. The products were filtered through diatomaceous earth, washing with dichloromethane. The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO₂, 50% to 60% ethyl acetate in *isohexane* as eluant) to give the title compound as a solid (100mg).

MS: APCI(+ve) 452 ($M+H^+$).

m.p. 172-177°C.

¹H NMR (400 MHz, d₆-DMSO) δ 8.07 (1H, d), 7.98 (1H, s), 7.66 (1H, d), 7.53 (1H, d), 7.47 (1H, d), 7.30 (1H, dd), 6.31 (1H, t), 3.20 (2H, d), 2.40 (3H, s), 2.22 (3H, s), 2.02 (3H, s), 1.74 (3H, d), 1.65 (3H, d), 1.57-1.61 (6H, m).

5

Pharmacological Analysis


Certain compounds such as benzoylbenzoyl adenosine triphosphate (bbATP) are known to be agonists of the P2X₇ receptor, effecting the formation of pores in the plasma membrane (Drug Development Research (1996), 37(3), p.126). Consequently, when the receptor is
10 activated using bbATP in the presence of ethidium bromide (a fluorescent DNA probe), an increase in the fluorescence of intracellular DNA-bound ethidium bromide is observed. The increase in fluorescence can be used as a measure of P2X₇ receptor activation and therefore to quantify the effect of a compound on the P2X₇ receptor.

15 In this manner, each of the title compounds of the Examples was tested for antagonist activity at the P2X₇ receptor. Thus, the test was performed in 96-well flat bottomed microtitre plates, the wells being filled with 250 µl of test solution comprising 200 µl of a suspension of THP-1 cells (2.5 x 10⁶ cells/ml) containing 10⁻⁴M ethidium bromide, 25 µl of a high potassium buffer solution containing 10⁻⁵M bbATP, and 25 µl of the high
20 potassium buffer solution containing concentrations of test compound typically from 30 µM – 0.001 µM. The plate was covered with a plastics sheet and incubated at 37 °C for one hour. The plate was then read in a Perkin-Elmer fluorescent plate reader, excitation 520 nm, emission 595 nm, slit widths: Ex 15 nm, Em 20 nm. For the purposes of comparison, bbATP (a P2X₇ receptor agonist) and pyridoxal 5-phosphate (a P2X₇ receptor
25 antagonist) were used separately in the test as controls. From the readings obtained, a pIC₅₀ figure was calculated for each test compound, this figure being the negative logarithm of the concentration of test compound necessary to reduce the bbATP agonist activity by 50%. Each of the compounds of the Examples demonstrated antagonist activity, having a pIC₅₀ figure > 5.5. For example, the following table shows the pIC₅₀
30 figures for a representative selection of compounds:

Compound of Example No.	pIC ₅₀
1	6.5
11	6.8

CLAIMS

1. A compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof,




wherein m represents 1, 2 or 3;


each R¹ independently represents a hydrogen atom or a halogen;

A represents C(O)NH or NHC(O);


Ar¹ represents a group



(II)




(III)



(IV)

or



(V)

one of R² and R³ represents halogen, nitro, NR⁴R⁵, hydroxyl, or a group

selected from (i) C₁-C₆ alkyl optionally substituted by at least one halogen and (ii)

C₁-C₆ alkoxy optionally substituted by at least one halogen, and the other of R² and R³ represents a hydrogen atom, halogen or a C₁-C₆ alkyl group optionally substituted by at least one halogen;

R⁴ and R⁵ each independently represent a hydrogen atom or a group selected from

C₁-C₆ alkyl and C₁-C₆ alkoxy, which C₁-C₆ alkyl or C₁-C₆ alkoxy group can be optionally substituted with at least one substituent selected from halogen and hydroxyl;

Ar² represents phenyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, which phenyl or

heteroaromatic ring is substituted by at least one substituent selected from CO_2R^6 , $\text{MC}_{1-6}\text{alkylCO}_2\text{R}^7$, $\text{C}_{1-6}\text{alkylsulphonylaminocarbonyl}$, NHR^8 , R^9 , XR^{10} , C(O)NHOH and $\text{NR}^{28}\text{R}^{29}$;

and which phenyl or heteroaromatic ring can further be optionally substituted by at least one substituent selected from halogen, nitro, $\text{NR}^{11}\text{R}^{12}$, hydroxyl, $\text{S(O)}_p\text{R}^{13}$, a $\text{C}_1\text{-C}_6$ alkoxy group which $\text{C}_1\text{-C}_6$ alkoxy group can be optionally substituted by at least one halogen, and a $\text{C}_1\text{-C}_6$ alkyl group which $\text{C}_1\text{-C}_6$ alkyl group can be optionally substituted by at least one substituent selected from halogen, hydroxyl, $\text{NR}^{14}\text{R}^{15}$, $\text{SO}_2\text{NR}^{16}\text{R}^{17}$, $\text{NR}^{18}\text{SO}_2\text{R}^{19}$, NHCOR^{20} and $\text{CONR}^{21}\text{R}^{22}$;

R^6 and R^7 each independently represent a hydrogen atom or a $\text{C}_1\text{-C}_6$ alkyl group;

R^8 represents CN , $\text{C}_1\text{-C}_6\text{alkylsulphonyl}$, $\text{C}_1\text{-C}_6\text{alkylcarbonyl}$, $\text{C}_1\text{-C}_6\text{alkoxycarbonyl}$, $\text{C}_1\text{-C}_6\text{alkylaminosulphonyl}$, or $(\text{di})\text{-C}_1\text{-C}_6\text{alkylaminosulphonyl}$;

R^9 and R^{10} each independently represent tetrazolyl or a 5- to 6-membered heterocyclic ring comprising from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulphur, which heterocyclic ring is substituted by at least one substituent selected from hydroxyl, $=\text{O}$ and $=\text{S}$;

M represents a bond, oxygen, S(O)_q or NR^{23} ;

X represents oxygen, S(O)_s , NR^{24} , $\text{C}_1\text{-C}_6\text{alkylene}$, $\text{O(CH}_2\text{)}_{1-6}$, $\text{NR}^{25}(\text{CH}_2)_{1-6}$, or $\text{S(O)}_t(\text{CH}_2)_{1-6}$;

p , q , s and t each independently represent 0, 1 or 2;

R^{28} and R^{29} together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring, which heterocyclic ring is substituted with at least one substituent independently selected from CO_2R^6 , $\text{MC}_{1-6}\text{alkylCO}_2\text{R}^7$, $\text{C}_{1-6}\text{alkylsulphonylaminocarbonyl}$, C(O)NHOH , NHR^8 , R^9 and XR^{10} , and which 3- to 8-

membered saturated heterocyclic ring can further be optionally substituted by at least one substituent independently selected from hydroxyl, halogen, $\text{C}_1\text{-C}_6$ alkoxy optionally substituted by at least one halogen, and a $\text{C}_1\text{-C}_6$ alkyl group which $\text{C}_1\text{-C}_6$ alkyl group can be optionally substituted by at least one substituent independently selected from halogen and hydroxyl; and

R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} and R^{25} each independently represent a hydrogen atom or a group selected from $\text{C}_1\text{-C}_6$ alkyl and $\text{C}_1\text{-C}_6$ alkoxy, which

C₁-C₆ alkyl or C₁-C₆ alkoxy group can be optionally substituted with at least one substituent selected from halogen and hydroxyl;

provided that:


- when m is 1 and Ar¹ is a group (II) and Ar² is phenyl substituted by XR¹⁰ in a position para to Ar¹ and X is CH₂, then R¹⁰ is not a 2,4-dioxothiazolyl group, and
- when m is 1 and Ar¹ is a group (II) and Ar² is phenyl substituted by MC₁.C₆alkylCO₂R⁷ in a position para to Ar¹, then M does not represent a bond.

2. A compound according to claim 1, wherein A represents NHC(O).

3. A compound according to claim 1 or claim 2, wherein Ar² represents phenyl, thienyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 nitrogen atoms.

4. A compound according to any one of claims 1 to 3, wherein Ar² is substituted by at least one substituent selected from carboxyl, -C₁.C₆alkylCO₂H, -OC₁.C₆alkylCO₂H, -N(C₁.C₆alkyl)C₁.C₆alkylCO₂H, -NHCN, -NHCO₂C₁.C₆alkyl, tetrazolyl and -OC₁.C₆alkyltetrazolyl, and wherein Ar² can further be optionally substituted by at least one substituent selected from halogen and C₁-C₆ alkyl.

5. A compound according to any one of claims 1 to 4, wherein Ar¹ represents a group



wherein R² represents halogen, nitro, NH₂, hydroxyl, or a C₁-C₆ alkyl optionally substituted by at least one halogen.

6. A compound according to claim 1 which is selected from:-

4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-4-carboxylic acid,

5 4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-3-carboxylic acid,

4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

2-Chloro-5-[6-(cyanoamino)pyrazinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
10 benzamide,

2-Chloro-5-[3-(cyanamino)pyrazinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

3-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyrazinecarboxylic acid,

15 3-[5-Chloro-4'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-2-pyridinyl]-benzoic acid,

2-Chloro-5-[3-[(methylsulfonyl)amino]pyrazinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(1-H-tetrazol-5-yl)pyrazinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
20 benzamide,

2-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid,

5-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid,

25 2-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-pyridinecarboxylic acid,

2-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-6-methyl-3-pyridinecarboxylic acid,

(2*S*)-2-[[[4'-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-
30 biphenyl]-2-yl]oxy]-propanoic acid,

[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-acetic acid,

3-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid,

5 5-Chloro-2-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid,

4'-Chloro-6-methyl-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

10 3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-thiophenecarboxylic acid,

6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid,

3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid,

15 2-Chloro-5-[2-(1*H*-tetrazol-5-yl)-3-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-oxazolecarboxylic acid,

20 4'-Chloro-4-methyl-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

6-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-*N*-(methylsulfonyl)-2-pyridinecarboxamide,

N-[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-glycine,

25 2-Chloro-5-[6-[(methylsulfonyl)amino]-2-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

[[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]oxy]-acetic acid,

30 2-Chloro-5-[3-(1*H*-tetrazol-5-ylmethoxy)-2-pyridinyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

4'-Chloro-4-methoxy-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

4-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-1-methyl-1*H*-pyrazole-3-carboxylic acid,

4-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-1-methyl-1*H*-pyrazole-5-carboxylic acid,

N-[3-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]pyrazinyl]-*N*-methyl-glycine ,

1-[3-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]pyrazinyl]-4-piperidinecarboxylic acid,

4'-Chloro-6-fluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

4'-Chloro-5-fluoro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-acetic acid,

[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-3-yl]oxy]-acetic acid,

(2*R*)-2-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-2-yl]oxy]-propanoic acid,

[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-4-yl]oxy]-acetic acid,

(2*S*)-2-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-3-yl]oxy]-propanoic acid,

4,4'-Dichloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

(2*S*)-2-[[4'-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl][1,1'-biphenyl]-4-yl]oxy]-propanoic acid,

3-Chloro-6-[4-chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid,

3-[4-Chloro-3'-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-pyridinecarboxylic acid,

[[2-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinyl]oxy]-acetic acid,

N-[2-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinyl]-glycine,

5 4'-Chloro-4,5-difluoro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

4'-Chloro-3'-[[2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

10 3-[4-Chloro-3-[[2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid,

4'-Chloro-4-fluoro-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

2-[5-Chloro-4-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-2-pyridinyl]-benzoic acid,

15 2-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-methyl-3-pyridinecarboxylic acid,

6-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-[(2-hydroxyethyl)methylamino]-3-pyridinecarboxylic acid,

20 3-[4-Methyl-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid,

4-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-1,3-dimethyl-1*H*-pyrazole-5-carboxylic acid,

2-[4-Methyl-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid,

25 3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridineacetic acid,

1-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-4-piperidinecarboxylic acid,

1-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-L-proline,

30 1-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-3-piperidinecarboxylic acid,

1-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-3-azetidinecarboxylic acid,

3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-6-methyl-2-pyridinecarboxylic acid,

5 3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-5-methyl-2-pyridinecarboxylic acid,

1-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-4-hydroxy-4-piperidinecarboxylic acid,

1-[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-5-fluoro-2-pyridinyl]-4-piperidinecarboxylic acid,

10 4'-Methyl-3'-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-[1,1'-biphenyl]-2-carboxylic acid,

1-[3-[4-Methyl-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinyl]-4-piperidinecarboxylic acid,

15 6-[4-Methyl-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-pyridinecarboxylic acid,

4-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-pyridinecarboxylic acid,

6-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-3-methyl-2-pyridinecarboxylic acid,


20 6-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-(trifluoromethyl)-2-pyridinecarboxylic acid, or

5-[6-(Acetylamino)-2-methyl-3-pyridinyl]-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide


25 or a pharmaceutically acceptable salt or solvate thereof.

7. A process for the preparation of a compound of formula (I) as defined in claim 1, or a pharmaceutically acceptable salt or solvate thereof, which comprises:-

30 (a) reacting a compound of formula




with a compound of formula



5 wherein one of Y and Z represents a displaceable group such as a metallic, organometallic or organosilicon group and the other of Y and Z represents a leaving group such as a halogeno or sulphonyloxy group and R^1 , m, A, Ar^2 , R^2 and R^3 are as defined for formula (I); or


10 (b) when Ar^2 is substituted by carboxyl, reacting a compound of formula (VI)-(IX) as defined in (a) above with a compound of formula



15 wherein Z is as defined in formula (X), and Ar^{2a} represents phenyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, followed by reaction with a base, then optionally followed by reaction with an acid; or

(c) when R^9 represents tetrazolyl, reacting a compound of formula (VI)-(IX) as defined in (a) above with a compound of formula (XI) as defined in (b) above, followed by reaction with a suitable source of azide; or


- 5 (d) when R^8 represents CN, C_{1-6} alkylsulphonyl, C_{1-6} alkylcarbonyl, C_{1-6} alkoxy carbonyl, C_{1-6} alkylaminosulphonyl, or (di)- C_{1-6} alkylaminosulphonyl, reacting a compound of formula (VI) – (IX) as defined in (a) above with a compound of formula



10

wherein L^1 represents a leaving group such as a halogeno or sulphonyloxy group, Ar^{2b} represents phenyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, and Z is as defined in formula (X), followed by reaction with a compound of formula

15



wherein V represents a hydrogen or a metallic group; or

- 20 (e) when Ar^2 is substituted by carboxyl, reacting a compound of formula (VI) - (IX) as defined in (a) above with a compound of formula (XII) as defined in (d) above, followed by reaction with a suitable source of cyanide, followed by reaction with a base, then optionally followed by reaction with an acid; or


- 25 (f) when R^9 represents tetrazolyl, reacting a compound of formula (VI)-(IX) as defined in (a) above with a compound of formula (XII) as defined in (d) above, followed by reaction with a suitable source of cyanide, followed by reaction with a suitable source of azide; or

- (g) when Ar^2 is substituted by carboxyl, reacting a compound of formula (VI)-(IX) as
30 defined in (a) above with a compound of formula (XII) as defined in (d) above, followed


by reaction with carbon monoxide and an alcohol in the presence of a suitable catalyst, followed by reaction with a base; or

(h) when Ar^2 represents a group of formula

5



reacting a compound of formula




10

with a suitable cyclodehydrating reagent followed by reaction with a suitable oxidising reagent followed by reaction with a base; or

(i) when M represents oxygen or NR^{23} , reacting a compound of formula (VI) – (IX) as

15


defined in (a) above, with a compound of formula (XII) as defined in (d) above, followed by reaction with a compound of formula




wherein M represents oxygen or NR^{23} , and R^{23} and R^7 are as defined in formula (I), optionally followed by reaction with a suitable base or acid; or

5 (j) when M represents oxygen or NR^{23} , reacting a compound of formula (XXI) as defined in (i) above, with a compound of formula (XII) as defined in (d) above, followed by reaction with a compound of formula (VI)-(IX) as defined in (a) above, optionally followed by reaction with a suitable base or acid; or

10 (k) when M represents oxygen or NR^{23} , reacting a compound of formula (VI)-(IX) as defined in (a) above, with a compound of formula




15 wherein Ar^{2c} represents phenyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, Z is as defined in formula (X), and M represents oxygen or NR^{23} , wherein R^{23} is as defined in formula (I), followed by reaction with either β -propiolactone or a compound of formula




wherein R^7 is as defined in formula (I), and L^1 is as defined in formula (XII), optionally followed by reaction with a suitable base or acid; or

25 (l) when X represents $\text{O}(\text{CH}_2)_{1-6}$ or $\text{NR}^{25}(\text{CH}_2)_{1-6}$ and R^{10} represents tetrazolyl, reacting a compound of formula (VI)-(IX) as defined in (a) above, with a compound of formula




30 wherein M^1 represents oxygen or NR^{25} , R^{25} is as defined in formula (I), Ar^{2d} represents a phenyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms

independently selected from nitrogen, oxygen and sulphur, and Z is as defined in formula (X), followed by reaction with a compound of formula




wherein L^1 is as defined in formula (XII), followed by reaction with a suitable source of azide; or

(m) when Ar^2 is substituted by carboxyl, reacting a compound of formula (VI)-(IX) as defined in (a) above with a compound of formula




wherein Z is as defined in formula (X), and Ar^{2e} represents a phenyl or 5- or 6-membered heteroaromatic ring comprising from 1 to 2 heteroatoms independently selected from nitrogen, oxygen and sulphur, followed by reaction with a suitable oxidising agent; or


(n) reacting a compound of formula




(XXIX),



(XXX),




(XXXI), or




(XXXII)

with a compound of formula



wherein one of R^{30} and R^{31} represents NH_2 and the other of R^{30} and R^{31} represents CO_2H ,
 5 $COBr$ or $COCl$, and Ar^2 , R^1 , R^2 , R^3 , R^6 and m are as defined in formula (I); or

(o) when R^{28} and R^{29} together with the nitrogen to which they are attached form a 3- to 8-
 membered saturated heterocyclic ring, which heterocyclic ring is substituted by CO_2R^6 ,
 reacting a compound of formula (VI) – (IX) as defined in (a) above, with a compound of
 10 formula (XII) as defined in (d) above, followed by reaction with a compound of formula



wherein R^6 , R^{28} and R^{29} are as defined in formula (I), optionally followed by reaction with
 15 a suitable base or suitable acid; or

(p) when R^{28} and R^{29} together with the nitrogen to which they are attached form a 3- to 8-
 membered saturated heterocyclic ring, which heterocyclic ring is substituted by CO_2R^6 ,
 reacting a compound of formula (XII) as defined in (d) above with a compound of formula
 20 (XXXIV) as defined in (o) above, followed by reaction with a compound of formula (VI) –
 (IX) as defined in (a) above, optionally followed by reaction with a suitable base or acid;

and optionally after (a), (b), (c), (d), (e), (f), (g), (h), (i), (j), (k), (l), (m), (n), (o) or (p)
 carrying out one or more of the following:

- converting the compound to a further compound of the invention


- forming a pharmaceutically acceptable salt or solvate of the compound.

8. A pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 6 in association with a pharmaceutically acceptable adjuvant, diluent or carrier.


9. A process for the preparation of a pharmaceutical composition as claimed in claim 8 which comprises mixing a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as defined in any one of claims 1 to 6 with a pharmaceutically acceptable adjuvant, diluent or carrier.

10. A compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 6 for use in therapy.

11. Use of a compound of formula (IA), or a pharmaceutically acceptable salt or solvate thereof, in the manufacture of a medicament for use in the treatment of an inflammatory disorder



wherein m represents 1, 2 or 3;
 each R¹ independently represents a hydrogen atom or a halogen;
 A represents C(O)NH or NHC(O);
 Ar¹ represents a group



one of R^2 and R^3 represents halogen, nitro, NR^4R^5 , hydroxyl, or a group
 5 selected from (i) C_1 - C_6 alkyl optionally substituted by at least one halogen and (ii)
 C_1 - C_6 alkoxy optionally substituted by at least one halogen, and the other of R^2 and R^3
 represents a hydrogen atom, halogen or a C_1 - C_6 alkyl group optionally substituted by at
 least one halogen;

R^4 and R^5 each independently represent a hydrogen atom or a group selected from

10 C_1 - C_6 alkyl and C_1 - C_6 alkoxy, which C_1 - C_6 alkyl or C_1 - C_6 alkoxy group can be optionally
 substituted with at least one substituent selected from halogen and hydroxyl;

Ar^2 represents phenyl or a 5- or 6-membered heteroaromatic ring comprising from 1 to 2
 heteroatoms independently selected from nitrogen, oxygen and sulphur, which phenyl or
 heteroaromatic ring is substituted by at least one substituent selected from CO_2R^6 , $MC_1.C_6$
 15 $alkylCO_2R^7$, C_{1-6} alkylsulphonylaminocarbonyl, NHR^8 , R^9 , XR^{10} , $C(O)NHOH$ and
 $NR^{28}R^{29}$;

and which phenyl or heteroaromatic ring can further be optionally substituted by at least
 one substituent selected from halogen, nitro, $NR^{11}R^{12}$, hydroxyl, $S(O)_pR^{13}$, a C_1 - C_6 alkoxy
 group which C_1 - C_6 alkoxy group can be optionally substituted by a halogen, and a

20 C_1 - C_6 alkyl group which C_1 - C_6 alkyl group can be optionally substituted by at least one
 substituent selected from halogen, hydroxyl, $NR^{14}R^{15}$, $SO_2NR^{16}R^{17}$, $NR^{18}SO_2R^{19}$,
 $NHCOR^{20}$ and $CONR^{21}R^{22}$;

R^6 and R^7 each independently represent a hydrogen atom or a $C_1.C_6$ alkyl group;

R^8 represents CN , $C_1.C_6$ alkylsulphonyl, $C_1.C_6$ alkylcarbonyl, $C_1.C_6$ alkoxy carbonyl, $C_1.C_6$
 25 alkylaminosulphonyl or (di)- $C_1.C_6$ alkylaminosulphonyl;

R^9 and R^{10} each independently represent tetrazolyl or a 5- to 6-membered heterocyclic ring
 comprising from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and

sulphur, which heterocyclic ring is substituted by at least one substituent selected from hydroxyl, =O and =S;

M represents a bond, oxygen, S(O)_q or NR²³;

X represents oxygen, S(O)_s, NR²⁴, C₁-C₆ alkylene, O(CH₂)₁₋₆, NR²⁵(CH₂)₁₋₆, or

5 S(O)_t(CH₂)₁₋₆;

p, q, s and t each independently represent 0, 1 or 2;

R²⁸ and R²⁹ together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring, which heterocyclic ring is substituted with at least one substituent independently selected from CO₂R⁶, MC₁-C₆ alkylCO₂R⁷, C₁-C₆

10 alkylsulphonylaminocarbonyl, C(O)NHOH, NHR⁸, R⁹ and XR¹⁰, and which 3- to 8-

membered saturated heterocyclic ring can further be optionally substituted by at least one substituent independently selected from hydroxyl, halogen, C₁-C₆ alkoxy optionally substituted by at least one halogen and a C₁-C₆ alkyl group which C₁-C₆ alkyl group can be optionally substituted by at least one substituent independently selected from halogen and

15 hydroxyl; and
R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴ and R²⁵ each independently represent a hydrogen atom or a group selected from C₁-C₆ alkyl and C₁-C₆ alkoxy, which C₁-C₆ alkyl or C₁-C₆ alkoxy group can be optionally substituted with at least one substituent selected from halogen and hydroxyl.

20 12. Use according to claim 11, wherein the inflammatory disorder is rheumatoid arthritis.

13. Use according to claim 11, wherein the inflammatory disorder is osteoarthritis.

25 14. Use according to claim 11, wherein the inflammatory disorder is asthma or chronic obstructive pulmonary disease.

15. Use of a compound of formula (IA), or a pharmaceutically acceptable salt or solvate thereof, in the manufacture of a medicament for use in the treatment of atherosclerosis.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2005/001251

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: see extra sheet

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL, WPI DATA, PAJ, CHEM ABS DATA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 03042190 A1 (PFIZER PRODUCTS INC.), 22 May 2003 (22.05.2003), page 1, line 31 - page 2, line 22; page 17, line 23 - page 22, line 28; examples 1-6; claims 1-10 --	1-15
X	WO 0061569 A1 (ASTRAZENECA AB), 19 October 2000 (19.10.2000), examples 1-88; claims 1-16 --	1-15
X	WO 0194338 A1 (ASTRAZENECA AB), 13 December 2001 (13.12.2001), examples 1-12; claims 1-16 --	1-15
A	WO 03080579 A1 (ASTRAZENECA AB), 2 October 2003 (02.10.2003), examples 1-208; claims 1-20 --	1-15

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

31 October 2005

Date of mailing of the international search report

01-11-2005

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Eva Johansson/EÖ

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2005/001251

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2004058270 A1 (PFIZER PRODUCTS INC.), 15 July 2004 (15.07.2004), example 32; claims 1-15 --	1-15
A	WO 9929661 A1 (ASTRA AKTIEBOLAG), 17 June 1999 (17.06.1999), examples 1-80; claims 1-12 --	1-15
A	WO 0144170 A1 (ASTRAZENECA AB), 21 June 2001 (21.06.2001), examples 1-76; claims 1-22 -- -----	1-15

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE2005/001251

C07D241/26, C07D241/14, C07D241/12, C07D213/24, C07D213/04,
C07D403/04, C07D401/04, C07D333/32, C07D327/04, C07D231/14,
C07C233/65, A61K31/192, A61K31/166, A61K31/4965, A61K31/497,
A61K31/435, A61P9/10, A61P11/00, A61P19/02, A61P37/00

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/10/2005

International application No.

PCT/SE 2005/001251

WO	03042190	A1	22/05/2003	AT	302196	T	15/09/2005
				BR	0204588	A	16/09/2003
				CA	2411544	A	12/05/2003
				DE	60205564	D	00/00/0000
				EP	1310493	A,B	14/05/2003
				JP	2003183263	A	03/07/2003
				MX	PA02009783	A	21/05/2003
				US	6927219	B	09/08/2005
				US	20030144293	A	31/07/2003

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/10/2005

International application No.

PCT/SE 2005/001251

WO	0061569	A1	19/10/2000	AU	774526	B	01/07/2004
				AU	3994700	A	14/11/2000
				AU	5547000	A	02/01/2001
				BR	0009651	A	08/01/2002
				CA	2368829	A	19/10/2000
				CN	1147483	C	28/04/2004
				CN	1353702	A,T	12/06/2002
				CZ	20013608	A	15/05/2002
				EE	200100525	A	16/12/2002
				EP	1171432	A	16/01/2002
				GB	0002330	D	00/00/0000
				HU	0202214	A	28/10/2002
				IL	145505	D	00/00/0000
				JP	2002541249	T	03/12/2002
				NO	20014894	A	10/12/2001
				NZ	514477	A	29/04/2003
				PL	350907	A	10/02/2003
				SK	13422001	A	09/05/2002
				TR	200102911	T	00/00/0000
				US	6492355	B	10/12/2002
				AP	200102041	D	00/00/0000
				AT	250036	T	15/10/2003
				AU	751103	B	08/08/2002
				AU	4950499	A	07/02/2000
				BR	9912109	A	02/05/2001
				CA	2336968	A,C	27/01/2000
				DE	69911415	D,T	08/07/2004
				DK	1095021	T	24/11/2003
				EE	200100010	A	17/06/2002
				EP	1095021	A,B	02/05/2001
				SE	1095021	T3	
				HK	1035715	A	00/00/0000
				HR	20010039	A	31/12/2001
				HU	0103224	A	28/01/2002
				IL	140346	D	00/00/0000
				JP	2002520395	T	09/07/2002
				NO	20010211	A	15/03/2001
				NZ	508923	A	27/09/2002
				PL	345388	A	17/12/2001
				SE	9901270	D	00/00/0000
				UA	64791	C	15/03/2001
				ZA	200108265	A	08/01/2003
<hr/>							
WO	0194338	A1	13/12/2001	AU	6447401	A	17/12/2001
				EP	1292590	A	19/03/2003
				GB	0013737	D	00/00/0000
				JP	2004506611	T	04/03/2004
				US	20030187031	A	02/10/2003
<hr/>							

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/10/2005

International application No.

PCT/SE 2005/001251

WO	03080579	A1	02/10/2003	AU	2003216013	A	00/00/0000
				EP	1399103	A	24/03/2004
				EP	1490341	A	29/12/2004
				JP	2004527322	T	09/09/2004
				SE	0200920	D	00/00/0000
				US	20040134103	A	15/07/2004
				US	20050090524	A	28/04/2005

WO	2004058270	A1	15/07/2004	AU	2003285716	A	00/00/0000
				NL	1025148	A	01/07/2004
				US	20040180894	A	16/09/2004

WO	9929661	A1	17/06/1999	AT	224360	T	15/10/2002
				AU	744280	B	21/02/2002
				AU	1791399	A	28/06/1999
				BR	9813390	A	03/10/2000
				CA	2312420	A	17/06/1999
				CN	1284057	A,T	14/02/2001
				DE	69808130	D,T	28/05/2003
				DK	1036059	T	02/12/2002
				EE	200000378	A	17/12/2001
				EP	1036059	A,B	20/09/2000
				SE	1036059	T3	
				ES	2184352	T	01/04/2003
				HU	0004434	A	28/05/2001
				ID	28003	A	00/00/0000
				IL	136369	D	00/00/0000
				JP	2001525392	T	11/12/2001
				NO	20002786	A	31/07/2000
				NZ	504447	A	26/04/2002
				PL	340906	A	12/03/2001
				PT	1036059	T	28/02/2003
				RU	2214997	C	27/10/2003
				SE	9704544	D	00/00/0000
				SI	1036059	T	28/02/2003
				SK	8432000	A	11/12/2000
				TR	200001605	T	00/00/0000
				US	6201024	B	13/03/2001
				US	6258838	B	10/07/2001
				US	6303659	B	16/10/2001
				US	20010003121	A	07/06/2001

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/10/2005

International application No.

PCT/SE 2005/001251

WO	0144170	A1	21/06/2001	AT	261933	T	15/04/2004
				AU	780506	B	24/03/2005
				AU	2244401	A	25/06/2001
				AU	2005202321	A	23/06/2005
				BR	0016395	A	27/08/2002
				CA	2393352	A	21/06/2001
				CN	1434794	A,T	06/08/2003
				CZ	20022093	A	15/01/2003
				DE	60009147	D,T	21/04/2005
				DK	1242364	T	21/06/2004
				EE	200200330	A	15/10/2003
				EP	1242364	A,B	25/09/2002
				SE	1242364	T3	
				EP	1352895	A	15/10/2003
				EP	1352896	A	15/10/2003
				EP	1352897	A	15/10/2003
				ES	2215777	T	16/10/2004
				GB	0015744	D	00/00/0000
				HK	1046678	A	03/12/2004
				HU	0300616	A	28/07/2003
				IL	150124	D	00/00/0000
				JP	2003517035	T	20/05/2003
				MX	PA02005789	A	18/09/2002
				NO	20022856	A	16/08/2002
				NZ	519378	A	27/02/2004
				PL	355913	A	31/05/2004
				PT	1242364	T	30/07/2004
				SI	1242364	T	31/08/2004
				SK	8412002	A	04/02/2003
				TR	200401432	T	23/08/2004
				US	6881754	B	19/04/2005
				US	20030013704	A	16/01/2003
				US	20050049303	A	03/03/2005
				GB	0017942	D	00/00/0000
				SE	9904651	D	00/00/0000
				ZA	200204125	A	25/08/2003
